Geological Survey of Canada
Commission géologique du Canada

BULLETIN 364

# SILURIAN AND DEVONIAN SPORE ZONES OF THE OLD RED SANDSTONE CONTINENT AND ADJACENT REGIONS

J.B. RICHARDSON
D.C. McGREGOR



1986

Canada

86-04937/

GEOLOGICAL SURVEY OF CANADA
BULLETIN 364

# SILURIAN AND DEVONIAN SPORE ZONES OF THE OLD RED SANDSTONE CONTINENT AND ADJACENT REGIONS

J.B. RICHARDSON
D.C. McGREGOR

1986

© Minister of Supply and Services Canada 1986

Available in Canada through

authorized bookstore agents and other bookstores

or by mail from

Canadian Government Publishing Centre
Supply and Services Canada
Ottawa, Canada K1A 0S9

and from

Geological Survey of Canada offices:

601 Booth Street
Ottawa, Canada K1A 0E8

3303-33rd Street N.W.,
Calgary, Alberta T2L 2A7

A deposit copy of this publication is also available
for reference in public libraries across Canada

Cat. No. M42-364E Canada: $10.00
ISBN 0-660-12112-3 Other countries: $12.00

Price subject to change without notice

**Critical readers**

*A.W. Norris*
*M. Streel*

**Scientific editor**

*N.C. Ollerenshaw*

**Editor**

*J. MacGillivray*

**Technical editor and layout**

*L. Machan-Gorham*

**Word processing**

*P. Greener*
*H. King*
*M. Varalta*

Artwork by the CARTOGRAPHY UNITS of
the Geological Survey of Canada,
Ottawa and Calgary

**Authors' addresses**

*J.B. Richardson*
*Department of Palaeontology*
*British Museum (Natural History)*
*Cromwell Road*
*London SW7 5BD, England*

*D.C. McGregor*
*Geological Survey of Canada*
*601 Booth Street*
*Ottawa, Ontario K1A 0E8*

*Original manuscript submitted: 83-08-29*
*Approved for publication: 85-08-27*

**Cover photos**

Left: **Retispora lepidophyta** (Kedo) Playford, enlarged X1000, from the Harrylock Formation, Hook Head, Ireland. It is a near-worldwide zonal indicator of Late Famennian and Strunian age (see p. 22-23). GSC type no. 73310; GSC loc. 99211.

Right: **Emphanisporites annulatus** McGregor, enlarged X1000, from the Stooping River Formation, Moose River Basin, Ontario. It is of near-worldwide occurrence in rocks of Emsian age (see p. 12-14). GSC type no. 41759; GSC plant loc. 8095.

## PREFACE

Correlation of rock units is important to the understanding of geological relationships. Spores are among the most useful fossils for this purpose, as evidenced by the fact that they have been valuable tools for hydrocarbon exploration in Canada and abroad for many years.

Spores are widespread in marine and nonmarine rocks of Silurian and Devonian age. However, to date no comprehensive spore zonation for the whole of the Silurian and Devonian systems has been proposed. This report describes such a zonation for North America and Europe. New spore zones are calibrated with zones already established for graptolites, conodonts, ammonoids and tentaculites found exclusively in marine strata. Thus, a basis for direct time correlation of widespread marine and continental rocks of the Silurian and Devonian of Canada is established in this report. It also provides data that may be useful for the international definition of geological boundaries.

*R.A. Price*
Director General
Geological Survey of Canada

## PRÉFACE

La corrélation des unités rocheuses tient un rôle important relatif à l'interprétation géologique. Parmi des fossils qui jouent un rôle majeur à cette fin, les spores prennent une place capitale, fait qui a été déjà mis en évidence au Canada et ailleurs dans l'exploration des hydrocarbures.

Des spores sont répandues dans les roches marines et non-marines du Silurien et du Dévonien. Néanmoins, jusqu'à présent, l'on n'a pas proposé une zonation compréhensive de spores pour l'ensemble des systèmes Silurien et Dévonien. Le présent rapport décrit une telle zonation pour l'Amérique du Nord et l'Europe. De nouvelles zones de spores s'étalonnent aux zones déjà établies de graptolites, de conodontes, d'ammonoïdes, et de tentaculites trouvés dans des strates exclusivement marines. Dès lors, une base pour la corrélation temporelle directe de roches marines et continentales très répandues du Silurien et du Dévonien du Canada se voit établie dans ce rapport. De plus, il présente des données d'utilité potentielle à la définition internationale de limites géologiques.

Le directeur général de la
Commission géologique du Canada
*R.A. Price*

## CONTENTS


1 Abstract/Résumé
1 Introduction
2 Acknowledgments
3 Zonal concept
4 Nomenclature of the zones
4 Criteria for characterizing the zones
6 Description of the zones
6 1. *Ambitisporites avitus* - *A. dilutus* Assemblage Zone
7 2. *Archaeozonotriletes chulus* var. *chulus* - *A. chulus* var. *nanus* Assemblage Zone
8 3. "*Emphanisporites*" cf. *protophanus* - cf. *Synorisporites verrucatus* Assemblage Zone
8 4. *Synorisporites libycus* - ?*Lophozonotriletes poecilomorphus* Assemblage Zone
9 5. *Synorisporites tripapillatus* - *Apiculiretusispora spicula* Assemblage Zone
10 6. *Emphanisporites micrornatus* - *Streelispora newportensis* Assemblage Zone
11 7. *Breconisporites breconensis* - *Emphanisporites zavallatus* Assemblage Zone
11 8. *Verrucosisporites polygonalis* - *Dictyotriletes emsiensis* Assemblage Zone
12 9. *Emphanisporites annulatus* - *Camarozonotriletes sextantii* Assemblage Zone
14 10. *Grandispora douglastownense* - *Ancyrospora eurypterota* Assemblage Zone
15 11. *Calyptosporites velatus* - *Rhabdosporites langii* Assemblage Zone
15 12. *Densosporites devonicus* - *Grandispora naumovii* Assemblage Zone
16 13. *Geminospora lemurata* - *Cymbosporites magnificus* Assemblage Zone
17 14. *Contagisporites optivus* var. *optivus* - *Cristatisporites triangulatus* Assemblage Zone
18 15. *Archaeoperisaccus ovalis* - *Verrucosisporites bulliferus* Assemblage Zone
20 16. *Auroraspora torquata* - *Grandispora gracilis* Assemblage Zone
21 17. *Rugospora flexuosa* - *Grandispora cornuta* Assemblage Zone
22 18. *Vallatisporites pusillites* - *Retispora lepidophyta* Assemblage Zone
23 19. *Verrucosisporites nitidus* - *Vallatisporites verrucosus* Assemblage Zone
24 References
36 Appendix


### Illustrations

#### Figures


in pocket 1. Approximate correlation of selected spore zones or assemblages with the assemblage zones described in this paper.
5 2. Stratigraphic ranges of nominal species of the spore assemblage zones, and comparison of the zones with the floral zones of Banks (1980) and with selected "events" in the evolution of spore morphology.
in pocket 3. Stratigraphic ranges of the characterizing species of the spore assemblage zones.
in pocket 4. Suggested approximate correlation of the spore assemblage zones with standard graptolite, conodont, and ammonoid zones, and with selected parts of stratigraphic successions in the U.K., France, Belgium, West Germany, North Africa, the United States and Canada that contain the zonal reference sections designated herein.
7 5. Geographical location of reference sections.
19 6. Approximate correlation of selected late Givetian and early Frasnian spore-bearing strata of the European U.S.S.R., Western Europe, and Arctic Canada.
20 7. Approximate correlation of the *torquata-gracilis/flexuosa-cornuta* boundary in New York State, based on spores and invertebrate faunas.


#### Plates


37 1-21. Illustrations of spores.


#### Tables


6 1. Conformity of spore zone circumscriptions with criteria 3 to 5 for definition of zones.
in pocket 2. Data for illustrated specimens.
in pocket 3. Lithology, depositional environment and fossil content of spore zones in the vicinity of the reference sections.

# SILURIAN AND DEVONIAN SPORE ZONES OF THE OLD RED SANDSTONE CONTINENT AND ADJACENT REGIONS


***Abstract***

Silurian and Devonian stratigraphic sections from marine and nonmarine strata of the Old Red Sandstone Continent and adjacent basins were studied palynologically and collated. Nineteen spore zones are proposed, some of which may be subdivided regionally. The zones are regarded as assemblage biozones. Each zone is based on the following: two characterizing and widely-occurring species after which the zone is named, at least one of which begins at the base of the zone; an association of other species of spores that typically occur together; and, for some zones, a major morphological "event", such as the first appearance of proximal radial muri (early Gorstian) and the first appearance of bifurcate-tipped spines (latest Emsian). For the bases of most of the zones, reference sections have been selected.

The wide geographical spread of the zones, especially those in the Lower Devonian, testifies to the apparently near cosmopolitan distribution of at least some components of the vegetation that produced the spores. On the other hand, in parts of the Upper Devonian, some spores (and therefore the plants that produced them) show a distinctly regional pattern of distribution. Most of the spore "events" do not coincide temporally with "events" that have been selected independently to delimit Siluro-Devonian plant megafossil zones.

***Résumé***

Des coupes stratigraphiques du Silurien et du Dévonien provenant de couches marines et non marines du continent des vieux grès rouges et de bassins voisins ont été étudiées d'un point de vue palynologique et elles ont été rassemblées. Dix-neuf zones de spores sont proposées, quelques-unes d'entre elles pouvant être subdivisées dans certaines régions. Ces zones sont considérées comme des biozones d'assemblage. Chaque zone est fondée sur 1) deux espèces caractéristiques à large distribution d'après lesquelles la zone est dénommée, au moins une des deux apparaissant à la base de la zone, 2) un assemblage d'autres espèces de spores qui sont typiquement présentes en même temps; et pour quelques zones 3) un "évènement" morphologique majeur tel que la première apparition de muri radiaux proximaux (Gorstein ancien) et la première apparition d'épines à extrémité bifurquée (Emsien le plus tardif). Des coupes de référence avaient été sélectionnées pour les limites inférieures de la plupart des zones.

La grande distribution géographique des zones, en particulier celles du Dévonien inférieur, atteste une distribution pratiquement cosmopolite d'au moins quelques éléments de la végétation qui a produit les spores. D'autre part, au Dévonien supérieur, quelques spores (et, par conséquent, les plantes qui les ont produites) montrent un schéma de distribution régional bien distinct. La plupart des "évènements" remarqués chez les spores ne coincïdent pas dans le temps avec des "évènements" qu'avaient été choisis indépendamment à délimiter les zones de mégafossiles végétaux siluro-dévoniens.


## INTRODUCTION

Spores with tetrad marks, produced by terrestrial plants, occur abundantly and in considerable variety in Late Silurian (Gorstian) and younger rocks, and in lesser numbers and diversity in strata at least as old as middle Llandovery. Trilete spores have been reported repeatedly from Llandovery rocks, and many have been dated convincingly using marine fossils. A number of authors have described spores and tetrads that they claim are of pre-Llandovery age (e.g. Combaz, 1967, Tremadoc; Timofeev, 1966, Cambrian, Early, Middle and Late Ordovician) but, in these works, either the age or the precise nature of the fossils is in dispute (some may be acritarchs), or the specimens are not adequately described or illustrated.

More recently, Gray, Massa and Boucot (1982) published excellent photographs of "permanent" tetrads, and *Ambitisporites*-like miospores broken out of tetrads, from rocks of supposed Caradoc (Late Ordovician) age. Their assemblage resembles those from the Llandovery of north Africa. In Llandovery rocks, single spores are well known, if sometimes rare, but there is a tendency for the spores to

remain in tetrads to a greater extent than in younger rocks. Thus, based on morphology alone, the "Caradoc" spores reported by Gray et al. (1982) could be Silurian. On the stratigraphic evidence presented so far by Gray et al. (1982) and Massa (1980), one must exercise caution in accepting a Caradoc age for them. However, "permanent" miospore tetrads have also been recorded from the Ordovician by Vavrdova (1982, Pl. 1, fig. 1, Pl. 4), from the Kosov Formation, which corresponds approximately to the upper Ashgillian *Glyptograptus bohemicus* Zone.

Fossil spores commonly are divided into two broadly defined types based on size. Megaspores, comprising morphographic species with a mean size of 200 µm or greater, appear in the late Siegenian, become progressively more abundant through the Emsian and Middle Devonian, achieve a remarkable differentiation in the Frasnian, and persist to the present day. Miospores, less than 200 µm in diameter, are first seen in rocks at least as old as the Early Silurian and are the most nearly ubiquitous of identifiable land plant remains in sediments of that age and younger. They occur in various continental facies and, significantly for biostratigraphy, in certain marine facies as well.

Some generalizations may be made regarding the nature of spore assemblages that occur in the principal, broadly categorized sedimentary environments, based on those recovered from Devonian sediments (e.g. Nazarenko, 1974) and from younger deposits (e.g. Stanley, 1969; Tschudy, 1969). The most abundant and diverse assemblages, and the largest and best preserved spores, are found in continental and neritic facies, with the greatest concentration and diversity being in the inner neritic zone but (except at deltas) some distance offshore. Spores may even occur in deep water sediments far from shore, but there they commonly are small, more limited in number and diversity, and less well preserved.

At present there is no internationally accepted spore zonation for the whole of the Silurian and Devonian. This report proposes such a zonation, based mainly on miospores, for the Old Red Sandstone (ORS) Continent and adjacent regions. With some modification, the scheme incorporates zones proposed for parts of the Devonian by Streel (1966, and later papers), Richardson (1974), McGregor and Camfield (1976) and others (see Figure 1 and McGregor, 1979b, figures 2-7), and considerable unpublished information, some of which is included in theses by Ahmed (1978), Ford (1972), Hassan (1982), Karathanasopoulos (1975), Riegel (1975) and Wood (1978). The data are drawn mainly from the authors' work on western European, North African and North American sequences, and from the literature on spores of the European U.S.S.R. An outline of this scheme was presented to the Carboniferous Congress in Moscow in 1975. Comments made at that meeting and subsequently by Soviet palynologists, have been considered in the selection of characteristic species, although the final result is solely the authors' responsibility. Correlation with spore assemblages of the Soviet Union is difficult, largely because Soviet and non-Soviet palynologists use different, independently-developed schemes of taxonomy and nomenclature.

The study of Silurian and Devonian spores is advancing rapidly. Considering the dynamic state of the subject, the zones we are proposing should not be regarded as final. More information is needed regarding the geographical distribution of the zones both within and, especially, beyond the ORS Continent, as well as about the relationship of the scheme to floral provinciality. For example, although little has been known about Devonian spores from China until recently, contemporary Chinese research suggests that some of our zones may be applicable there (see Figure 1). Furthermore, some of the key species and morphological "events" in the spore succession can be recognized in sequences from South America, Antarctica and Australia, and are thus almost worldwide. The present emphasis on the zonation of the ORS Continent and adjacent sedimentary basins, therefore, is not so much a result of known biogeographical constraints as it is of the fact that most of the thoroughly described and precisely dated assemblages are from those regions.

We expect that as work progresses it will be possible to recognize additional zones, as has been true, for example, with conodont zonation. Indeed, this is already possible on an intra-regional basis (see Figure 1). In addition, as more data accumulate, it will be possible to test more fully whether the zones are in fact synchronous over the whole ORS region, or possibly even beyond this region.

The International Union of Geological Sciences Subcommission on Devonian Stratigraphy has defined the bases of the Eifelian, Frasnian and Famennian stages as at the lower boundaries of the *Polygnathus costatus partitus*, Lower *Polygnathus asymmetricus* and Middle *Palmatolepis triangularis* conodont zones, respectively. We accept these recommendations. For those stages with bases that are as yet not defined internationally, we have related the spore zones provisionally to faunally-defined versions of the relevant stage boundaries (see Figure 4).

The Subcommission on Devonian Stratigraphy has also recommended Lochkovian, Pragian and Emsian as the international standard stage nomenclature for the Lower Devonian. As a result of this decision, the Rhenish stages Gedinnian and Siegenian assume the status of regional stages. Although we acknowledge the wisdom of this decision as it applies to faunal correlations, we prefer to retain the Rhenish terminology at this time, as correlations of the spore-bearing Rhenish facies with Lochkovian and Pragian strata of the pelagic facies still are not securely established.

Unless stated to the contrary, all of the proposed zones and spore features discussed on the following pages are based on spores with tetrad marks, i.e. spores thought to have been produced by land plants. Citations of spore nomenclature, including "?" and "cf." designations, are given in the form in which they were originally published. Information on the provenance and repository of the specimens illustrated herein is given in Table 2 and the Appendix.

**Acknowledgments**

W. Riegel of the Geologisches Institut, Göttingen, and M. Streel of the University of Liège are especially thanked for their significant contributions to the improvement of the manuscript, Riegel in the early stages and Streel in the final stages of its preparation. We are also grateful to A. Arkhangelskaya and T.V. Byvsheva of the All-Union Petroleum Scientific Research Institute for Geological Exploration (VNIGNI), Moscow; E.W. Bamber, A.W. Norris and T.T. Uyeno of the Geological Survey of Canada, Calgary; E. Turnau of the Institute of Geological Sciences, Kraków, and C.G. Adams of the British Museum (Natural History) for valuable comments; to G.S. Nowlan of the Geological Survey of Canada, Ottawa, for comments on Figure 4; to Gao Lianda of the Academy of Geological Sciences, Beijing, W. Riegel and M. Streel for unpublished information; to K. Higgs of the Geological Survey of Ireland, Dublin, P. van Veen of the

Norwegian Petroleum Directorate, Stavanger, B. Owens of the Institute of Geological Sciences, Keyworth, England, A. Traverse of Pennsylvania State University, W. Riegel and M. Streel for providing photographs of spores; and to John Wells of Cornell University and Irving Tesmer of Buffalo College, New York State, for help on the stratigraphy of New York State and northern Pennsylvania.

## ZONAL CONCEPT

Much has been written about the structure, definition, and limitations of biozones (e.g. Teichert, 1958; Hancock, 1977; Holland et al., 1978). However, the theoretical basis for the creation of palynological zones has received relatively little publicity, especially amongst proponents of the traditional, marine, faunal orthochronologies. It therefore seems appropriate to explain the rationale underlying the zonal scheme we are proposing.

Among the most significant characteristics of Recent and fossil spores are their abundance, their relative resistance to destruction, and the extraordinary scope of their dispersal by wind and especially by water. Assemblages of dispersed spores may be preserved in both continental and marine sediments. Except in autochthonous coals, each assemblage as a whole tends to reflect the combined vegetation of various parts of the drainage basin, whereas individual components of the assemblage may vary both qualitatively and quantitatively according to sedimentary factors and local concentrations of the parent vegetation onshore. Assemblages are commonly composed of thirty or more stratigraphically useful species, and many thousands of specimens may be present in a few grams of rock.

Widespread, apparently synchronous "first appearances" of certain taxa and form features of spores occur repeatedly in the geological column, and the Silurian and Devonian are not exceptional in this respect. Such occurrences are assumed to reflect evolution, and morphological antecedents can be suggested for many taxa. Whatever their evolutionary antecedents, certain (but not all) species and structural features seem to have attained widespread geographical distribution instantaneously in terms of geological time. Consequently, we have accorded considerable biochronological significance to first appearances of selected species and form features (Fig. 2). Upper stratigraphical limits (disappearances) may have been affected by such factors as persistence of relict floras, or redeposition, and so are given less weight in the definition of zonal boundaries.

On the other hand, there are some species that apparently do not make their first appearance contemporaneously in the sections that have been studied. A case can be made for significant ecological and hydrodynamic control, in addition to the large-scale geographical controls mentioned below, as an explanation for the seemingly heterochronous appearance of some species. Many authors have discussed this phenomenon. For example, Streel (1967), Richardson (1969) and McGregor (1981) have dealt with several species that occur in the vicinity of the Middle-Upper Devonian boundary. This phenomenon emphasizes the advisability of comparing whole assemblages comprising shared, characteristic taxa for long-distance correlation using spores.

Most of the zones defined here are, to some degree, geographical composites. That is, the lists of characteristic species commonly include certain elements of contemporaneous assemblages that appear to be restricted geographically. *Archaeoperisaccus* spp. in the Frasnian, and *Cornispora* spp. in the Famennian are examples of species with well-defined geographical limits (see McGregor, 1979b). However, each zone has shared, characteristic taxa that permit interregional temporal correlation of the zone.

The zones we are proposing are characterized by species that may have different inherent values for long-distance time-correlation. Some species do not have global distribution but may characterize floral provinces or realms, whereas others may represent more local variations that nevertheless have to be taken into account for interregional correlation.

Recently, some workers have adopted the use of palynological "phases" to facilitate regional correlation (Van der Zwan, 1980; Loboziak and Streel, 1981). Phases are considered to represent successive, evolutionary(?) steps in the gradual compositional development of palynological assemblages (Van der Zwan, 1980, p. 192). The phase concept may be useful, but it needs to be defined more explicitly and distinguished clearly from local and regional zonation.

In view of the foregoing, we believe the most reasonable basis for defining spore zones that can be correlated on an intercontinental scale is a combination of the co-occurrence of whole assemblages comprising several characteristic taxa, and the first appearance of selected species and form features. The earliest records of one or two selected species, observed in a variety of facies and irrespective of relative abundance (which may, however, be significant for local correlations), are used to define the base of each zone in a reference section. Assemblages occurring in consistent order of succession, each characterized by several species, provide the basis for correlations on a global scale. The Oppel zone, as defined in the International Stratigraphic Guide (Hedberg, 1976), which is one variant of the assemblage zone of the recently revised North American Stratigraphic Code (1983, p. 863), closely approximates this approach. The "global zones" of Mamet (1977) are in some ways similar in concept, but differ in being based on supposed phyletic lineages. Johnson (1979) has presented a useful discussion of both of these zonal concepts, which he calls the "overlap range zone" and the "lineage zone". Our zones are not interval zones, as they are defined in part, and correlated, by assemblages of characteristic taxa.

Because we use first occurrence of species rather than maximum abundance or frequency of occurrence to define zone boundaries, some nominal species may have their acme above the zone bearing their name. An example is *Geminospora lemurata*, the acme of which is not in the *lemurata-magnificus* Zone but in the succeeding *optivus-triangulatus* Zone.

We have not attempted to establish spore lineages, although such attempts, based on morphological "successions", might be just as feasible for spores as for other fossils. The most secure basis for recognition of lineages might be the demonstrated relationship of certain spore forms to their parent plants (*vide inter alia* Gensel, 1979). Visscher (1971, p. 27) was of the opinion that the typological concept in palynological taxonomy may be fundamentally unsuited to the recognition of phyletic lineages. He suggested that the "deme" system, which emphasizes the gradation rather than the uniformity of morphological characters, might facilitate the recognition of evolutionary

lines. We suspect, however, that the proposal of empirically-based "phyletic" lineages on the data presently available could encourage entrenched thinking and consequent circular reasoning, and would not improve the precision of our correlations.

Silurian and Devonian plant megafossil zones (Banks, 1980) are in part reflected in the proposed spore zonation (Fig. 2). For example, the diversification of land plants (base of Banks's zone II) and the appearance of a group of advanced progymnosperms (near the base of Banks's zone V) approximately correlate with the spore changes that occur at the beginning of our *micromatus-newportensis* and *lemurata-magnificus* zones. On the other hand, there are other megafossil zones, such as Banks's zone III, whose bases at present do not coincide with the base of any spore zone, although plants within zone III, for example *Dawsonites*, *Sawdonia*, and *Psilophyton*, have produced spores of the *Apiculiretusispora* type, which are most abundant in the upper Siegenian and Emsian (the range of Banks's zone). Some apparent discrepancies between the two zonal systems may stem from the difficulty of correlating continental strata (from which most plant megafossils have been obtained) with marine stages. Spore studies can make a major contribution to the solution of this problem. Other discrepancies between the spore and plant megafossil zonations may be harder to resolve, for the reasons given by Banks (1980, p. 3-5).

## NOMENCLATURE OF THE ZONES

The name given to each spore zone comprises (a) the names of two widely-occurring and characteristic taxa, referred to as "nominal species", and (b) a statement of the kind of zone. On its first citation, the name of the zone is given in full, for example the *Contagisporites optivus* var. *optivus-Cristatisporites triangulatus* Assemblage Zone. In subsequent citations, the name of the zone is abbreviated – the *optivus-triangulatus* Zone.

The use of combinations of letters derived from the nominal species, e.g. "OT Zone" for the *optivus-triangulatus* Zone, is not recommended. Such letter designations are hard to remember, convey no information as to the composition of the zone, and may be confused with or duplicated by previously or subsequently established two-letter zonal designations, such as those of Clayton et al. (1977). When two-letter designations are also used for subzones (Clayton et al., 1978), phases (Van der Zwan, 1980), and acritarch zones (Loboziak et al., 1983), even greater confusion is possible.

If the names of the nominal taxa change, the name of the zone should be changed accordingly.

The numbers preceding the headings of the zone descriptions in the text refer to Figure 5, and are not intended to be part of the nomenclature of the zones.

## CRITERIA FOR CHARACTERIZING THE ZONES

1. At least one of the nominal species selected for each zone begins at the base of the zone in the reference section. Most of the nominal species are geographically widespread.

2. The zones are recognized by the presence of a substantial number of the characteristic species. They are not circumscribed by the local or total ranges of the nominal species. A nominal species may also occur in stratigraphically higher zones, or may even be absent from its zone in some places. The known stratigraphic ranges of the nominal species and the other characteristic species are given in figures 2 and 3.

3. The bases of some zones are also characterized by the first manifestations of major features of spore morphology, such as the appearance of bifurcate-tipped processes in the late Emsian at the base of the *douglastownense-eurypterota* Zone. We refer to these phenomena as "events" (Fig. 2). Such palynological "events" may transgress conventional taxonomic divisions. They are of greater temporal and geographical magnitude than the "events" referred to by Hughes (1975, p. 142) in relation to his "biorecords".

4. A good zonal base exists if the defining "events" and the first occurrence of characteristic taxa are not accompanied by an observable lithological or structural discontinuity, or associated with mass extinction or renewal of taxa, and if a similar succession of taxa and "events" has been observed in two or more sedimentary basins.

5. The base of each zone is defined with respect to a specified stratigraphic section in which the zone is characteristically developed, and which also contains a zonally significant fauna. The top of each zone is defined by the base of the succeeding zone.

The standards described in the foregoing paragraphs represent ideal conditions for the definition of spore zones. Some of the zones described here do not fit all the established criteria (see Table 1). For example, the bases of the *avitus-dilutus* and *chulus-nanus* zones have not been defined in reference sections. Nevertheless, after some hesitation, we have included these two zones because, whatever their exact limits, they appear to be consistent in composition over a wide geographical area, and they represent the oldest known assemblages of trilete spores.

***Figure 2.*** *Stratigraphic ranges of nominal species of the spore assemblage zones, and comparison of the zones with the floral zones of Banks (1980) and with selected "events" in the evolution of spore morphology. Broken horizontal lines indicate imprecisely defined zonal limits. Broken vertical lines indicate uncertain age and/or identification of species. The dotted line indicates uncertain correlation between the top of the Downtonian and the base of the Gedinnian.*


SERIES
STAGE
SPORE ZONE
1. Ambitisporites avitus
2. Ambitisporites dilutus
3. Archaeozonotriletes chulus var. chulus
4. Archaeozonotriletes chulus var. nanus
5. "Emphanisporites" cf. protophanus
6. cf. Synorisporites verrucatus
7. Synorisporites libycus
8. ?Laphozonotriletes poeciomorphus
9. Synorisporites tripapillatus
10. Apiculiretusispora spicula
11. Emphanisporites micrornatus
12. Streelispora newportensis
13. Breconisporites breconensis
14. Emphanisporites zavallatus
15. Verrucosisporites polygonalis
16. Dictyotriletes emsiensis
17. Emphanisporites annulatus
18. Camarozonotriletes sextantii
19. Grandispora douglastownense
20. Ancyrospora eurypterota
21. Calyptosporites velatus
22. Rhabdosporites langii
23. Densosporites devonicus
24. Grandispora naumovii
25. Geminospora lemurata
26. Cymbosporites magnificus
27. Contagisporites optivus var. optivus
28. Cristatisporites triangulatus
29. Archaeoperisaccus ovalis
30. Verrucosisporites bulliferus
31. Auroraspora torquata
32. Grandispora gracilis
33. Rugospora flexuosa
34. Grandispora cornuta
35. Vallatisporites pusillites
36. Retispora lepidophyta
37. Verrucosisporites nitidus
38. Vallatisporites verrucosus
EVENT
FLORAL ZONE
Banks, 1980
UPPER DEVONIAN
TOURNAISIAN (part)
FAMENNIAN
FRASNIAN
MIDDLE DEVONIAN
GIVETIAN
EIFELIAN
LOWER DEVONIAN
EMSIAN
SIEGENIAN
GEDINNIAN
PRIDOLI
DOWNTONIAN
LUDLOW
LUDFORDIAN
GORSTIAN
WENLOCK
HOMERIAN
SHEINWOODIAN
LLANDOVERY
TELYCHIAN
AERONIAN
RHUDDANIAN
niltidus-verrucosus
pusillites-lepidophyta
flexuosa-cornuta
torquata-gracilis
ovalis-bulliferus
optivus-triangulatus
lemurata-magnificus
devonicus-naumovii
velatus-langii
douglastownense-eurypterota
annulatus-sextantii
polygonalis-emsiensis
breconensis-zavallatus
micrornatus-newportensis
tripapillatus-spicula
libycus-poeciomorphus
cf. protophanus-verrucatus
chulus-nanus
avitus-dilutus
Vacuolate zona
Reappearance of foveoreticulate pseudosaccus
Prominent verrucate patina
Multifurcate spines
Monolete + zona-pseudosaccus
Camerate with thick exoexine
Distally sculptured bizonate cingulum; irregular large verrucae; irregular foveoreticulate pseudosaccus
Bifurcate spines; distinct increase in spore size
Spinose zona-pseudosaccus
Monolete mark
Megaspores (>200 µm)
Undifferentiated cingulum
Biform sculpture
Bizonate cingulum
Sloughing sculptured exoexine
Zona
Interradial proximal papillae; reticulate sculpture
Apiculate sculpture (on trilete spores)
Foveolate sculpture
Proximal radial muri
Verrucate sculpture
Patina
First undoubted trilete mark
VII Rhacophyton
VI Archaeopteris
V Svalbardia
IV Hyenia
III Psilophyton
II Zosterophyllum
I Cooksonia

GSC

**TABLE 1**

Conformity of spore zone circumscriptions with criteria 3 to 5 for definition of zones

| ASSEMBLAGE ZONE | "EVENT" AT BASE | "GOOD" BASE | REFERENCE SECTION | FAUNAL CONTROL |
|---|---|---|---|---|
| *nitidus-verrucosus* | - | X | X | - |
| *pusillites-lepidophyta* | - | X | X | X |
| *flexuosa-cornuta* | - | X | X | - |
| *torquata-gracilis* | X | X | X | X |
| *ovalis-bulliferus* | - | X | X | X |
| *optivus-triangulatus* | - | X | X | - |
| *lemurata-magnificus* | - | X | X | - |
| *devonicus-naumovii* | X | X | X | X |
| *velatus-langii* | - | - | X | X |
| *douglastownense-eurypterota* | X | X | X | X |
| *annulatus-sextantii* | - | - | X | X |
| *polygonalis-emsiensis* | X | X | X | X |
| *breconensis-zavallatus* | X | X | X | X |
| *micrornatus-newportensis* | - | X | X | X |
| *tripapillatus-spicula* | X | - | X | X |
| *libycus-poecilomorphus* | X | X | X | X |
| cf. *protophanus-verrucatus* | X | X | X | X |
| *chulus-nanus* | X | X | - | X |
| *avitus-dilutus* | X | X | - | X |

The first occurrence of a nominal species in the reference section for the zone may not represent the earliest chronostratigraphic record of the species, because of environmental and sedimentological constraints on the movement of spores to the site of deposition. Nevertheless, the stratigraphic level at which the lowest-occurring nominal species appears in the reference section shall be the "anchoring point" for determining the species content of the zone, until redefinition of the zone becomes necessary.

Figure 4 shows the approximate correlation of the spore zones with standard graptolite, conodont, ammonoid and tentaculite zones, based on published and unpublished information from many sources. Figure 4 also gives the stratigraphic position of the reference sections designated herein for the bases of the spore zones. Figure 5 shows the geographical location of the reference sections. Table 3 summarizes the lithology, depositional environments and fossil content of the spore zones in the vicinity of the reference sections.

In the following description of the zones, the plate and figure numbers given in brackets following the names of the characteristic species refer to illustrations in this bulletin.

## DESCRIPTION OF THE ZONES

### 1. *Ambitisporites avitus* - *A. dilutus* Assemblage Zone

***Age.*** Early and early Late Llandovery (Aeronian and early Telychian).

***Reference section for the base of the zone.*** None.

***Characteristic species.***

*Ambitisporites avitus* Hoffmeister, 1959, Pl. 1, figs. 1-8 (Pl. 1, fig. 1).

*A. dilutus* (Hoffmeister) Richardson and Lister, 1969; Hoffmeister, 1959, Pl. 1, figs. 9-13 (Pl. 1, fig. 2).

*Retusotriletes warringtonii* Richardson and Lister, 1969, Pl. 37, figs. 7 and 8 (Pl. 1, fig. 3).

*Tetrahedraletes medinensis* Strother and Traverse, 1979, Pl. 1, figs. 14-17 (Pl. 1, figs. 4, 5).

***Description.*** Laevigate, equatorially crassitate, trilete miospores and laevigate and sculptured monads, dyads and "permanent" tetrads. Spores are mainly small (mostly 20-40 µm in diameter), subtriangular or subcircular.

***Distribution.*** Upper Pentamerus Beds and lower Hughley Shales, Shropshire (Richardson in Aldridge et al., 1979), approximately *Monograptus sedgwickii* and *M. turriculatus* graptolite zones (Hill, 1974); in subsurface, Middle Llandovery of Libya (Hoffmeister, 1959; age determination




***Figure 5.*** *Geographical location of reference sections. Numbers refer to numbered zones in text. The limits of the Old Red Sandstone Continent are taken from Anderton et al., 1979, with revisions. The continental assembly is provisional.*

by Berry in Gray and Boucot, 1971); lower Massanutten Sandstone, Virginia (Pratt et al., 1978); Tuscarora Formation, Pennsylvania (Strother and Traverse, 1979). The Medina Group, Rhuddanian Stage, Lower Llandovery, New York State (Miller and Eames, 1982) contains a rich assemblage of spore-like fossils, but does not contain the diagnostic species of this zone.

***Remarks.*** "Permanent" tetrads (*Tetrahedraletes*) and dyads (*Dyadospora* Strother and Traverse, 1979) may be present. Some tetrads and dyads are enclosed by a single diaphanous membrane. Sculptured, alete, spore-like fossils, for example *Strophomorpha* Miller and Eames (1982), have been described from New York State and Pennsylvania, where tetrads and dyads along with sculptured alete spore-like fossils (monads) may dominate the assemblage.

### 2. *Archaeozonotriletes chulus* var. *chulus* - *A. chulus* var. *nanus* Assemblage Zone

***Age.*** Approximately Late Llandovery to late Wenlock (Telychian to late Homerian).

***Reference section for the base of the zone.*** None.

***Characteristic species.***

*Archaeozonotriletes chulus* (Cramer) Richardson and Lister var. *chulus*; Richardson and Lister, 1969, Pl. 43, figs. 1-6 (Pl. 1, figs. 6-8).

*A. chulus* (Cramer) Richardson and Lister var. *nanus* Richardson and Lister, 1969, Pl. 43, figs. 10 and 11 (Pl. 1, fig. 9).

*Ambitisporites avitus* Hoffmeister; see previous zone.

*A. dilutus* (Hoffmeister) Richardson and Lister; see previous zone.

?*A.* cf. *divellomedium* Chibrikova 1959; Richardson and Lister, 1969, Pl. 43, fig. 12 (Pl. 1, fig. 10).

*Retusotriletes warringtonii* Richardson and Lister; see previous zone.

*Tetrahedraletes medinensis* Strother and Traverse; see previous zone.

**Description.** First appearance of trilete, patinate, laevigate miospores with a thin proximal surface that bears characteristic compression folds concentric with, and located just inside, the equatorial thickening. First appearance of subcircular, possibly alete spores with polar(?), proximal(?) circular diaphanous area (?*Archaeozonotriletes* cf. *divellomedium*) in the upper part of the zone. Persistence of species from the previous zone, except for many of the sculptured "spore-like microfossils" (see Strother and Traverse, 1979 and Miller and Eames, 1982).

**Distribution.** Upper Coalbrookdale Beds (*Cyrtograptus lundgreni* graptolite biozone), Shropshire, England (Richardson and Lister, 1969); subsurface of Libya (Richardson and Al-Ameri, unpublished). An assemblage from the Lettergesh Formation of Ireland (Smith, 1975), dated as early Wenlock from graptolite evidence, could belong to either the *chulus-nanus* or the *avitus-dilutus* zone, as it does not contain *Archaeozonotriletes chulus*.

**Remarks.** Palynomorphs that appear identical to ?*A.* cf. *divellomedium* have been described from marine Silurian rocks of Spain by Cramer (1966a) under the name *Saturnus laevigatus*, a form regarded by Cramer as *incertae sedis*. However, similar forms have been recovered from nonmarine strata, and therefore probably were derived from land plants (Richardson and Lister, 1969). This form ranges into the Lower Devonian.

### 3. "*Emphanisporites*" cf. *protophanus* - cf. *Synorisporites verrucatus* Assemblage Zone

**Age.** Late Wenlock and early Ludlow (late Homerian to early Gorstian).

**Provisional reference section for the base of the zone.** Type area of the Wenlock Series, Much Wenlock Limestone Formation, approximately 4 m above its base, Harton Hollow Wood Quarry, 213.4 m south of Harton Village, Shropshire, England (Richardson and Lister, 1969).*

**Range of the zone.** Much Wenlock Limestone Formation (*Monograptus ludensis* graptolite zone) to the lower part of the lower Elton Beds [*Neodiversograptus nilssoni* graptolite zone(?)].

**Characteristic species.**

"*Emphanisporites*" cf. *protophanus* Richardson and Ioannides, 1973, Pl. 2, fig. 7 (Pl. 1, fig. 12).

cf. *Synorisporites verrucatus* Richardson and Lister; Richardson and Ioannides, 1973, Pl. 6, figs. 17-21 (Pl. 1, figs. 11, 13, 14).

*Ambitisporites dilutus* (Hoffmeister) Richardson and Lister; see previous zone.

*Archaeozonotriletes chulus* (Cramer) Richardson and Lister (both varieties); see previous zone.

?*A.* cf. *divellomedium* Chibrikova; see previous zone.

*Retusotriletes abundo* Rodriguez, 1978, Pl. 1, fig. 6 (Pl. 1, figs. 15, 16).

**Description.** Appearance of patinate alete(?) forms with verrucate sculpture; persistence of trilete spores with an equatorial crassitude. Appearance of spores that are possibly alete and that have incipient proximal ribs ("*Emphanisporites*" cf. *protophanus*). Beginning of retusoid spores with distinctly invaginated curvaturae and circular or subcircular amb (*Retusotriletes abundo*) high in the zone. In comparison, *R. warringtonii* of the preceding zones has a subtriangular amb and equatorial curvaturate thickenings that rarely invaginate onto the proximal surface.

**Distribution.** Much Wenlock Limestone Formation and lower Elton Beds, Shropshire, England (Richardson and Lister, 1969); calcareous silts and nodular limestones, Much Wenlock Limestone Formation(?), Trostrey Wood, Monmouthshire (Richardson, unpublished); subsurface of Libya (Richardson and Ioannides, 1973; Richardson and Al-Ameri, unpublished); San Pedro Formation, northern Spain (Rodriguez, 1978).

**Remarks.** Alete spores with "distal" verrucate sculpture, similar to cf. *Synorisporites verrucatus*, were recorded by Cramer (1966a) as *Hispanaediscus verrucatus* (*incertae sedis*).

### 4. *Synorisporites libycus* - ?*Lophozonotriletes poecilomorphus* Assemblage Zone

**Age.** Gorstian and Ludfordian. Graptolite zones *Neodiversograptus nilssoni* to approximately *Neocucullograptus kozlowskii* (see Bassett et al., 1982).

*Recently a research student, Neil Burgess, working at the British Museum (Natural History), located "*Emphanisporites*" cf. *protophanus* and cf. *Synorisporites verrucatus* in the upper Coalbrookdale Formation below Harton Hollow Wood Quarry in the type Wenlock area. Discovery of these specimens necessitates lowering the base of the cf. *protophanus* - *verrucatus* Zone to at least the upper lower Homerian in the graptolite upper *lundgreni* Zone.

***Reference section for the base of the zone.*** British Petroleum borehole B2-34, sample 18, Ghadames Basin, northern Libya (Richardson and Ioannides, 1973). Graptolites in this sequence indicate a late Wenlock or early Ludlow age (*op. cit.*, p. 258). Accessory reference section: top of the lower Elton Beds, junction of Elton Lane with the track to Evenhay Farm, 800 m southeast of Elton village and 6.4 km southwest of Ludlow, Shropshire, England. Spores are rare in this section (Richardson and Lister, 1969). The accessory reference section was chosen because it is in the type area of the Ludlow Series, and because the primary reference section is in a borehole.

***Range of the zone.*** In Shropshire, upper part of the lower Elton Beds to the top of the upper Whitcliffe Beds.

***Characteristic species.***

*Synorisporites libycus* Richardson and Ioannides, 1973, Pl. 7, figs. 1-6 (Pl. 2, figs. 1-3).

*?Lophozonotriletes poecilomorphus* Richardson and Ioannides, 1973, Pl. 7, figs. 9-15 and Pl. 8, figs. 1, 4-6 (Pl. 2, figs. 6, 7).

cf. *Brochotriletes* sp. A, Richardson and Ioannides, 1973, Pl. 4, figs. 5-6, 8-10 (Pl. 2, figs. 8, 9).

*"Emphanisporites" protophanus* Richardson and Ioannides, 1973, Pl. 2, figs. 8-12 (Pl. 2, fig. 12).

"*E.*" cf. *protophanus* Richardson and Ioannides; see previous zone.

*E. neglectus* Vigran, 1964, Pl. 1, figs. 14-16 (Pl. 2, fig. 4).

cf. *Synorisporites downtonensis* Richardson and Lister, 1969, Pl. 40, fig. 6 (Pl. 2, fig. 11).

*Synorisporites* cf. *verrucatus* Richardson and Ioannides, 1973, Pl. 6, figs. 11-16 (Pl. 2, figs. 5, 10).

*Tetraletes variabilis* Cramer, 1966b, Pl. 1, figs. 7, 8, 10 (Pl. 2, fig. 16).

***Description.*** First appearance of proximal radial muri on trilete spores (*Emphanisporites*), distal murornate sculpture (*?Lophozonotriletes*, cf. *Synorisporites downtonensis*), and apiculate tetragonal tetrads (*Tetraletes*). Beginning of apiculate sculpture on trilete spores. Appearance of foveolate sculpture (cf. *Brochotriletes*). Persistence of all nominal species of the previous zones.

***Distribution and regional variation.*** Acacus Formation of Libya (Richardson and Ioannides, 1973). Middle Elton, Bringewood, Leintwardine and Whitcliffe beds of Shropshire, England (Richardson and Lister, 1969; Richardson, unpublished), based on the first appearance of murornate sculpture; the nominal species were not found. Assemblages described by Aristova and Arkhangelskaya (1976) from the Kuresaare Beds contain *Retusotriletes* with wholly proximal curvaturae, *Emphanisporites*, and possibly *Apiculiretusispora* and *Brochotriletes*. All of these elements occur for the first time in the *libycus-poecilomorphus* Zone. The nominal species of this zone occur in the upper Tarabuco Formation and the lower Santa Rosa Formation of the Tarabuco region, Bolivia (McGregor, 1984). Other possible occurrences of this zone are in the Mississinewa Shale of Indiana (Wood, 1978), and the lower member of the Barlow Inlet Formation of Cornwallis Island, Canada (McGregor and Narbonne, 1978; Thorsteinsson and Uyeno, 1980). In the Ludlow, *Brochotriletes* and similar forms are so far known from Libya (Richardson and Ioannides, 1973), Spain (Rodriguez, 1978), Estonia (Aristova and Arkhangelskaya, 1976), and Bolivia (McGregor, 1984).

***Remarks.*** When further work on the systematic description of upper Gorstian and Ludfordian spores is completed it may be possible to separate the upper part of the *libycus-poecilomorphus* Zone on the basis of *Apiculiretusispora* spp.

### 5. *Synorisporites tripapillatus - Apiculiretusispora spicula* Assemblage Zone

***Age.*** Downtonian, sensu White, 1950; approximately equivalent to the Pridolian (see Cocks et al., 1971, p. 127; Kaljo, 1978; Richardson et al., 1981). A detailed review of the Downton Series is given by Bassett et al. (1982).

***Reference section for the base of the zone.*** Ludford Lane, Ludlow, Shropshire, England, base of the Downton Castle Sandstone Group, immediately above the Ludlow Bone Bed (Richardson and Lister, 1969). In the type area, no spores were recovered from the Ludlow Bone Bed, which is lenslike and, although persistent in many sections in the Welsh Borderland, is absent to the west and southwest of the region. This and the other bone beds in the sequence do not appear to represent marked discontinuities (Martinsson, 1977, p. 49).

***Range of the zone.*** Downton Castle Sandstone Group, Shropshire. Many of the characteristic species have not been found in the Temeside and Ledbury formations, but study of these beds is still preliminary.

***Characteristic species.***

*Synorisporites tripapillatus* Richardson and Lister, 1969, Pl. 40, figs. 7-9 (Pl. 2, figs. 17, 18).

*Apiculiretusispora spicula* Richardson and Lister, 1969, Pl. 38, figs. 3, 4 (Pl. 2, fig. 14).

*Ambitisporites* sp. B of Richardson and Ioannides, 1973, Pl. 6, figs. 7-9 (Pl. 3, fig. 1).

*Amicosporites splendidus* Cramer, 1967, Pl. 2, figs. 31, 40, 41 (Pl. 2, fig. 13).

*Apiculiretusispora synorea* Richardson and Lister, 1969, Pl. 38, figs. 5, 6 (Pl. 2, fig. 15).

*Cymbosporites echinatus* Richardson and Lister, 1969, Pl. 42, figs. 1-5 (Pl. 3, fig. 2).

*C. verrucosus* Richardson and Lister, 1969, Pl. 42, figs. 6-8 (Pl. 3, figs. 4, 5).

*Emphanisporites neglectus* Vigran, 1964; see previous zone.

*E. splendens* (Richardson and Ioannides) Richardson and Ioannides 1979; 1973, Pl. 3, figs. 12-15; Pl. 4, figs. 1-4, 7 (Pl. 3, fig. 3).

*Synorisporites downtonensis* Richardson and Lister, 1969, Pl. 40, figs. 4. 5 (Pl. 3, figs. 8, 9).

*S. verrucatus* Richardson and Lister, 1969, Pl. 40, figs. 10-12 (Pl. 3, figs. 6, 7).

***Description.*** First appearance of interradial proximal papillae (*S. tripapillatus*; *Ambitisporites* sp. B, ? = *A. tripapillatus* Moreau-Benoit, 1976), distally sculptured patinate spores with thin contact areas (*Chelinospora* and *Cymbosporites*), and reticulate sculpture (*Chelinospora*). Persistence of nearly all species from the previous zone.

***Distribution and regional variation.*** Downton Castle Sandstone Group, England and Wales (Richardson and Lister, 1969); upper Acacus Formation, Libya (Richardson and Ioannides, 1973); upper beds of the Oued Tifist Formation, Algeria (Jardiné and Yapaudjian, 1968). The upper Acacus Formation and the Oued Tifist Formation have not been faunally dated.

The zone may also occur in the Kaugatuma and Okhesaare(?) beds of Estonia (Aristova and Arkhangelskaya, 1976), the Borshchov Horizon of Podolia (Arkhangelskaya, 1980a; Richardson et al., 1981), and the Santa Rosa Formation of the Tarabuco region of Bolivia (McGregor, 1984). It has not been confirmed in North America, but spores possibly of this zone, as yet unstudied, are present in strata of Pridolian age in Canada, for example the Clam Bank Formation of Newfoundland (Martin, 1978, p. 77; McGregor, unpublished), and the Devon Island Formation of Devon Island, Queen Elizabeth Islands (McGregor, unpublished).

*Emphanisporites rotatus* is present high in the Acacus Formation of Downtonian age (based on spore correlation) in Libya; in parts of the San Pedro Formation assigned to the Ludlow and Downtonian by Rodriguez (1978); and in the poudingue de Fépin (lower Gedinnian) and the Oignies Formation (mid-Gedinnian) of Belgium (Streel et al., 1981). It has not been found below the Senni Beds (Siegenian) in the Anglo-Welsh sequence or the basal Stooping River Formation (early Siegenian) in Canada. *Emphanisporites splendens* is so far only known from Spain, north Africa (Richardson, Rasul and Al-Ameri, 1981) and China (Gao, 1981).

***Remarks.*** Work in progress on Downtonian spores by one of us (JBR) may result in subdivision of this zone.

### 6. *Emphanisporites micrornatus* - *Streelispora newportensis* Assemblage Zone

***Age.*** Early (but not earliest) and early late Gedinnian, by indirect correlation (Richardson et al., 1981; Richardson, unpublished; Blieck, 1982). *Pteraspis leathensis* and *Rhinopteraspis crouchi* vertebrate zones.

The base of the Gedinnian in the type area is considered to be, in this paper, approximately equivalent to the Siluro-Devonian boundary. Bultynck (1982) recorded conodonts of the *Icriodus woschmidti* Zone (latest Pridolian and earliest Lochkovian) in the boreholes at Artois, northern France, and also recorded the probable presence of this zone in the lower Gedinne Beds at Naux ("20 km southeast of Gedinne"). Palynological samples (Steemans, 1982a) from the Fépin Conglomerate and Haybes Beds, basal Gedinnian of the type locality (Lahonry Quarry), however, have yielded a *micrornatus-newportensis* assemblage that may be correlated with the Chortkov Beds of the Podolian sequence (Richardson et al., 1981). This assemblage is early but not earliest Devonian, and is younger than the *Monograptus uniformis* and *Icriodus woschmidti* zones. The Fépin Conglomerate is therefore regarded here as early but not earliest Gedinnian.

***Reference section for the base of the zone.*** Ross-Tewkesbury Spur Motorway (M. 50), Hereford and Worcester, England (Holland and Richardson, 1977; Richardson, unpublished). Base of the St. Maughan's Group, immediately above the prominent calcrete that "... is close to or encloses the Downtonian-Dittonian boundary and includes the "*Psammosteus*" Limestone ..." (Allen and Dineley, 1976, p. 13).

***Range of the zone.*** Lower and middle Ditton Group (base sensu White, 1950) Shropshire. The upper part of the zone is not well documented, because, so far, only poor assemblages have been found from the upper Ditton Group in the Clee Hills (Richardson, unpublished).

***Characteristic species.***

*Emphanisporites micrornatus* Richardson and Lister, 1969, Pl. 38, figs. 10, 11 (Pl. 4, fig. 1).

*Streelispora newportensis* (Chaloner and Streel) Richardson and Lister, 1969, Pl. 41, figs. 3-6 (Pl. 3, figs. 10, 11).

*Acinosporites salopiensis* Richardson and Lister, 1969, Pl. 39, figs. 5, 6 (Pl. 3, figs. 12, 13).

*Apiculiretusispora plicata* (Allen) Streel, 1967, Allen, 1965, Pl. 94, figs. 6-9 (Pl. 4, fig. 2).

*Chelinospora cassicula* Richardson and Lister, 1969, Pl. 42, figs. 10-12 (Pl. 4, figs. 6, 7).

*Cymbosporites proteus* McGregor and Camfield, 1976, Pl. 1, figs. 8-16, 21-23 (Pl. 4, figs. 3, 4).

*Emphanisporites epicautus* Richardson and Lister, 1969, Pl. 38, figs. 13-15 (Pl. 4, fig. 5).

*Perotrilites microbaculatus* Richardson and Lister, 1969, Pl. 39, figs. 7-13 (Pl. 4, figs. 8, 9).

*Retusotriletes maculatus* McGregor and Camfield, 1976, Pl. 1, fig. 6 (Pl. 4, fig. 10).

***Description.*** First appearance of prominent, flimsy zona (e.g. "spore zonée sp. A", Steemans, 1981), this feature rare until the succeeding zone. Beginning of sloughing, sculptured exoexine. Increase in the variety of proximally radially-ribbed spores (*Emphanisporites*), including the appearance and common occurrence of specimens with conate and granulate distal sculpture (*E. micrornatus*). Proliferation of reticulate sculpture (*Chelinospora* and, in the upper part of the zone,

*Dictyotriletes*). Persistence of spores with proximal, interradial papillae. Most spores are less than 40 µm in diameter.

***Distribution and regional variation.*** Ditton Group, Clee Hills, Shropshire, and St. Maughan's Group, Herefordshire, England; Red Marl Group, Breconshire, Wales (Richardson and Lister, 1969; Richardson, unpublished); Arbuthnott Group, Scotland (Richardson, 1967; Ford, 1972; Richardson et al., 1984); poudingue de Fépin and lower arkose d'Haybes (Steemans, 1982a) and Fooz Formation (Steemans, 1981), Dinant Synclinorium, Belgium; subsurface of Libya (Richardson et al., 1981; Richardson and Al-Ameri, unpublished); Strypa Horizon of the Dniestr Series (Shepeleva, 1963); the Chortkov and Ivane horizons of the Tiver series and the Ustetko Horizon of the lower Dniestr series (Arkhangelskaya, 1980a) of Podolia. *Emphanisporites micromatus* and other species consistent with the *micromatus-newportensis* Zone occur in the Kenogami River Formation of Northern Ontario (McGregor and Camfield, 1976). Some elements of the zone are present in the Massif Armoricain of France: the Saint Ceneré section (Moreau-Benoit, 1976) and the Saint-Pierre-sur-Erve section (D'Erceville, 1979). *S. newportensis*, but not *E. micromatus*, has been recorded from the Lower Devonian of Qujing, Yunnan Province, China (Gao Lianda, personal communication to JBR, 1981).

*E. micromatus* first appears 30 to 60 m above the base of the zone in the sections examined in the Brown Clee area, and is common in samples from the lower and middle Dittonian (Richardson, unpublished). It also appears in Podolia somewhat later than *S. newportensis*, in the Ivane Horizon (Richardson et al., 1981), and is not yet present in assemblages containing *S. newportensis* in the basal Gedinnian of the type region of the stage (Steemans, 1982a).

### 7. *Breconisporites breconensis* - *Emphanisporites zavallatus* Assemblage Zone

***Age.*** Late Gedinnian (Richardson et al., 1982). Approximately *Althaspis leachi* vertebrate zone.

***Reference section for the base of the zone.*** Belgium, southeastern Dinant Synclinorium, Nonceveux section along the road between Remouchamps and Trois-Ponts (Steemans, 1981, 1982b; Richardson et al., 1982); topographic map I.G.M., 1:25,000, Sheet 49, 3-4; Lambert grid X: 246500 and Y: 129750. The base of the zone is 35 m above the base of the Bois d'Ausse Formation. Accessory reference section, South Wales, Brecon Beacons, Glyn Tarell stream section (Hassan, 1982), map reference SN 9752 2163. The lowest strata known to be in the zone are close to the base of the Senni Beds, that is the lowest productive sample occurs within, but not at, the base of the basal sandstone unit of the Senni Beds (Richardson et al., 1982).

***Range of the zone.*** At Nonceveux, the zone includes at least 270 m of the Bois d'Ausse Formation, beginning at the lowest spore-bearing locality 35 m above the base of the Bois d'Ausse. In the southern part of the Dinant Basin the zone covers at least 400 m of strata. In the Black Mountains and Brecon Beacons, it covers the lower 64 to 90 m, respectively, of the Senni Beds except for the first 3 to 4 m of the basal sandstone unit. It is associated with the heterostracan *Althaspis leachi* (White) in Belgium (Richardson et al., 1982).

***Characteristic species.***

*Breconisporites breconensis* Richardson, Streel, Hassan and Steemans, 1982, Pl. 1, figs. 7-11 (Pl. 4, fig. 11).

*Emphanisporites zavallatus* Richardson, Streel, Hassan and Steemans, 1982, Pl. 1, figs. 1-6 (Pl. 4, fig. 13).

*Apiculiretusispora plicata* (Allen) Streel; see previous zone.

*Camptozonotriletes* sp. cf. *C. caperatus* McGregor; Hassan, 1982, Pl. 30, figs. 8, 9, 11 (Pl. 4, fig. 12).

*Cirratriradites* sp. A, Streel et al., 1981, Pl. 3, figs. 10, 11 (Pl. 4, fig. 14).

*Clivosispora verrucata* McGregor var. *verrucata*, McGregor, 1973, Pl. 7, figs. 4, 5, 10 (Pl. 4, figs. 15, 16).

***Description.*** First appearance of bizonate spores with distal annulate thickening (*Breconisporites breconensis*), coarsely sculptured *Emphanisporites* (*E. zavallatus*), and cingulate verrucate spores (*Clivosispora*). Appearance of zonate spores with verrucate or murornate sculpture (*Camptozonotriletes*).

***Distribution.*** Lower and middle Bois d'Ausse Formation, southeast part of Dinant Synclinorium, and upper Oignies and St. Hubert formations, southern Dinant Synclinorium in Belgium; lower and middle Senni Beds, Brecon Beacons and Black Mountains, Powys and Gwent, south Wales (Hassan, 1982); subsurface of the Moesian Platform, Romania (Beju, 1967); possibly in the Shyashuvian Suite, Lithuania (Arkhangelskaya, 1978).

### 8. *Verrucosisporites polygonalis* - *Dictyotriletes emsiensis* Assemblage Zone

***Age.*** Siegenian (and earliest Emsian?). Approximately *Rhinopteraspis dunensis* vertebrate zone, by indirect correlation with Blieck (1982). The precise relationships between the *polygonalis-emsiensis* Zone and the boundaries of the *dunensis* Zone are not known.

***Reference section for the base of the zone.*** Belgium, the southeastern Dinant Synclinorium, Nonceveux section along the road between Remouchamps and Trois-Ponts (Godefroid et al., 1982); topographic map I.G.M., 1:25,000, sheet 49, 3-4; Lambert grid X:248750 and Y:129750. The base of the zone is in the Bois d'Ausse Formation, 81 m below the base of the Solières Formation. Accessory reference section, middle Senni Beds, Craig Cerrig-gleisiad, Brecon Beacons, South Wales, map reference SN 9610 2200 (Richardson and Hassan, unpublished). The base of the zone is about 15.5 m above the base of the section. Samples from the underlying basal Senni Beds contain a *breconensis-zavallatus* assemblage.

***Range of the zone.*** Middle and upper Senni Beds; the overlying Brownstones Group has so far proved barren of spores.

***Characteristic species.***

*Verrucosisporites polygonalis* Lanninger, 1968, Pl. 22, fig. 19 (Pl. 5, figs. 1, 2).

*Dictyotriletes emsiensis* (Allen) McGregor 1973, Pl. 5, fig. 15 (Pl. 5, figs. 4, 5).

*Apiculiretusispora arenorugosa* McGregor, 1973, Pl. 2, figs. 15, 25 (Pl. 6, fig. 6).

*A.* cf. *brandtii* Streel, in Edwards, 1968, Pl. 132, figs. 8-10 (Pl. 5, fig. 9).

*A. plicata* (Allen) Streel; see previous two zones.

*Breconisporites breconensis* Richardson, Streel, Hassan and Steemans; see previous zone.

*Brochotriletes foveolatus*? Naumova; McGregor, 1973, Pl. 4, figs. 20, 23, 24 (Pl. 5, fig. 3).

*B. hudsonii* McGregor and Camfield, 1976, Pl. 3, figs. 1, 2 (Pl. 6, figs. 3, 4).

*Camptozonotriletes caperatus* McGregor, 1973, Pl. 7, figs. 1, 2 (Pl. 6, fig. 7).

*Clivosispora verrucata* McGregor var. *convoluta* McGregor and Camfield, 1976, Pl. 2, figs. 13-21 (Pl. 5, figs. 10, 11).

*Dibolisporites eifeliensis* (Lanninger) McGregor, 1973, Pl. 3, figs. 17-22, 26 (Pl. 5, fig. 6).

*D. wetteldorfensis* Lanninger; McGregor, 1973, Pl. 3, figs. 23-25 (Pl. 5, figs. 7, 8).

*D.* n. sp. K, Hassan, 1982, Pl. 12, figs. 1-6, 8-10 (Pl. 6, fig. 5).

*Dictyotriletes favosus* McGregor and Camfield, 1976, Pl. 2, figs. 5, 6 (Pl. 6, figs. 1, 2).

*Emphanisporites decoratus* Allen, 1965, Pl. 97, figs. 15-18 (Pl. 5, figs. 12, 13).

***Description.*** Appearance and proliferation of biform sculpture (*Dibolisporites*), including sculptural elements with expanded tips (*D. eifeliensis*). Proliferation of *Apiculiretusispora* spp., and diversification of *Dictyotriletes* spp. and coarsely sculptured *Emphanisporites*. Proximal interradial papillae are rare. First rare megaspores, *Dictyotriletes canadensis*, high in the zone (Grande Grève Formation, McGregor, unpublished).

***Distribution.*** Middle and upper Senni Beds, Brecon Beacons, Breconshire, and Mill Bay Formation (Cosheston Group), Dyfed, South Wales (Richardson, 1967; Richardson and Lister, 1969; Richardson and Thomas in Williams 1978); Grande Grève Formation, Gaspé (McGregor and Owens, 1966; McGregor, unpublished); Stooping River Formation, Northern Ontario (McGregor and Camfield, 1976); Upper Bois d'Ausse Formation and Solières, Acoz, Anor, St. Michel and Petigny formations, Dinant Basin, Belgium (Streel et al., 1981; Steemans, 1981, 1982b; Godefroid et al., 1982); southeast Massif Armoricain, France (Moreau-Benoit, 1976); and possibly the lower part of the Kemeri Series of Lithuania (Arkhangelskaya, 1980a), and the Snowblind Bay Formation of Cornwallis Island, Arctic Canada (McGregor, unpublished).

*Dictyotriletes emsiensis* appears below *Verrucosisporites polygonalis* in Ontario (McGregor and Camfield, 1976) and in Belgium (Streel, pers. comm., 1984) but above *V. polygonalis* in Gaspé (McGregor, 1977). Neither species has as yet been found in the Brecon Beacons. We have selected the first appearance of *V. polygonalis* to mark the base of the zone in the primary reference section.

### 9. *Emphanisporites annulatus* - *Camarozonotriletes sextantii* Assemblage Zone

***Age.*** Early and early late Emsian. Approximately *Polygnathus dehiscens* Zone to *Polygnathus serotinus* Zone inclusive, in the conodont zonation.

***Reference section for the base of the zone.*** Lower Singhofen Group, Beinhausen Formation, Gefell Member, southeast Eifel region (Fuchs, 1974; Riegel and Karathanasopoulos, 1982). Spores have not been found in the lowermost Emsian Ulmen Group, which lies immediately below the Gefell Member. Below the Ulmen Group, the Saxler Schichten (Herdorf Group) of uppermost Siegenian age contains spores, but not the nominal species or other spores typical of the *annulatus-sextantii* Zone, with the exception of *Apiculiretusispora brandtii*. A more suitable reference section, in stratigraphic continuity with underlying spore-bearing beds, may be found as more work is done on strata of this age.

***Range of the zone.*** Base of Beinhausen Formation (Gefell Member) to high Wetteldorf Formation.

***Characteristic species.***

*Emphanisporites annulatus* McGregor, 1961, Pl. 1, figs. 5, 6 (Pl. 6, figs. 9, 10).

*Camarozonotriletes sextantii* McGregor and Camfield, 1976, Pl. 4, 13, 14, 16-18 (Pl. 6, figs. 11, 12).

*Acinosporites lindlarensis* Riegel var. *lindlarensis*, Riegel, 1968, Pl. 19, figs. 11-16 (Pl. 6, fig. 8).

*Apiculatasporites microconus* (Richardson) McGregor and Camfield, 1982; Richardson, 1965, Pl. 89, fig. 3 (Pl. 7, fig. 1).

*Apiculiretusispora brandtii* Streel, 1964, Pl. 1, figs. 6-10 (Pl. 7, fig. 5).

*A. plicata* (Allen) Streel; see previous two zones.

*Calyptosporites biornatus* (Lanninger) Richardson, 1974; Lanninger, 1968, Pl. 25, fig. 12 (Pl. 7, fig. 4).

*Camptozonotriletes* cf. *aliquantus* Allen, 1965; Riegel, 1973, Pl. 15, figs. 4, 5 (Pl. 7, fig. 7).

*Dibolisporites echinaceus* (Eisenack) Richardson, 1965, Pl. 89, figs. 5, 6 (Pl. 7, figs. 2, 3).

*Dibolisporites eifeliensis* (Lanninger) McGregor; see previous zone.

*Dictyotriletes canadensis* McGregor, 1973, Pl. 5, figs. 21-23 (Pl. 7, fig. 8).

*D. gorgoneus* Cramer, 1967, Pl. 3, figs. 69, 72 (Pl. 7, fig. 6).

*Emphanisporites schultzii* McGregor, 1973, Pl. 6, fig. 14 (Pl. 7, fig. 9).

*Verruciretusispora dubia* (Eisenack) Richardson and Rasul, 1978, Pl. 1, fig. 6 (Pl. 7, fig. 10).

***Description.*** Greatest abundance and diversification of proximally radially ribbed spores (*Emphanisporites*). First appearance of distally annulate forms of *Emphanisporites*, for example *E. erraticus* and *E. annulatus*, the latter with an indistinctly delimited annulus in early Emsian and sharply delimited annulus in later Emsian. Diversification of biform sculpture (*Dibolisporites*). First appearance of *D. echinaceus* sensu Richardson (1965), *Apiculatasporites microconus* and *Acinosporites lindlarensis*, species that are more common in the succeeding zone. First prominent-spined zonate-pseudosaccates in the upper part of the zone. First rare monolete spores. Diversification of small megaspores.

***Distribution and regional variation.*** Beinhausen Formation, Gefell Member (Riegel and Karathanasopoulos, 1982), Stadtfeld, Klerf, Wiltz and Wetteldorf formations (Lanninger, 1968; Schultz, 1968), and Hunsrück Shale (Karathanasopoulos, 1975), in the Rheinisches Schiefergebirge; subsurface of southern England (Richardson and Rasul, 1978); "Emsian 2-3" of Sart-Tilman, Liège (Streel, 1967) and Wépion and Pesche beds in the Dinant Synclinorium (Streel et al., 1981) in Belgium; subsurface of Romania (Beju, 1967); Daleje Shale, Czechoslovakia (McGregor, 1979a); Ouan Kasa Formation and upper Tadrart Formation of Libya (Massa and Moreau-Benoit, 1976; Moreau-Benoit, 1979, 1980); York River Formation and lower Battery Point Formation of Gaspé (McGregor, 1977), La Garde Formation and Atholville beds of lower Restigouche River, Quebec and New Brunswick (Dineley and Williams, 1968), Sextant and Stooping River formations of Northern Ontario (McGregor and Camfield, 1976), and Vendom Fiord Formation of Ellesmere Island (McGregor, 1974 and unpublished) in Canada; Reuterskiøldfjellet Sandstone of Vestspitsbergen (Allen, 1965, 1967). Possibly in the Horlick Formation of Ohio Range, Antarctica (Kemp, 1972); La Vid Shales, Spain (Cramer, 1967); Danlin Formation, Kueichow, China (Gao and Hou, 1975); Malobachat Horizon of Salair, U.S.S.R. (Andreeva, 1973); and the upper part of the Kemeri Series of Lithuania (Arkhangelskaya, 1980a).

The upper part of this zone may occur in the Takata, Muradymovo, Vanyashkino and Vyazovo deposits of the western slope of the South Urals, where Chibrikova (1978, p. 111) has reported spores similar to the spores from the upper Emsian of the Rhine area, Belgium, Canada, and Spitsbergen. Spores with prominent, bifurcate-tipped spines, that typify the succeeding *douglastownense-eurypterota* Zone, including the upper Wetteldorf-Heisdorf-Lauch sequence of the Rhine region, have not been reported from these beds. According to Chibrikova (1972, p. 33; 1978), the Vanyashkino and Vyazovo are characterized by an influx of large, sculptured zonate-pseudosaccate spores (*Hymenozonotriletes* of the Soviet nomenclature). This event may correlate with the first appearance of similar spores in the upper part of the *annulatus-sextantii* Zone in western Europe and Canada (see Figure 2). The Vyazovo beds contain conodonts of the *Polygnathus inversus* and *P. serotinus* zones (Maslov et al., 1983), which is additional evidence that the Vyazovo deposits are older than the Heisdorf and uppermost Wetteldorf beds of the Eifel region (Weddige, 1977).

In Czechoslovakia, *Camarozonotriletes sextantii* is rare in the Daleje Shale near Srbsko, high in the tentaculite *Nowakia cancellata* Zone (McGregor, 1979a), but *E. annulatus* has not been found, perhaps owing to insufficient study.

The first large, prominent-spined, zonate-pseudosaccate spores appear well up in the *annulatus-sextantii* Zone, but they are not abundant until the succeeding *douglastownense-eurypterota* Zone. Such spores occur in the Wiltz Formation and possibly, if records of *Calyptosporites biornatus* are substantiated, as early as the Klerf Formation of the southeast Eifel region (Riegel and Karathanasopoulos, 1982); high in the *N. cancellata* Zone of the Barrandian region (McGregor, 1979a); and high in the Battery Point Formation of eastern Gaspé (McGregor, 1977).

*Camarozonotriletes sextantii* is rare in the Battery Point Formation of Gaspé, but is abundant in the Sextant and Stooping River formations of Northern Ontario. T.T. Uyeno comments as follows (personal communication to DCM, 1984): "In the Stooping River Formation it [*C. sextantii*] begins about 65 m above the apparent base of the conodont *dehiscens* Zone. This base cannot be located precisely, because the conodont faunas of the Stooping River Formation consist primarily of icriodontids, not zone-defining polygnathids. The Pragian-Zlichovian boundary occurs within the *dehiscens* Zone (Klapper and Johnson, 1980, p. 403), but as the lower limit of this zone cannot be located, the stadial boundary is similarly elusive." In Belgium, Steemans (1982b) found *C. sextantii* in the Petigny Formation, i.e. in the upper Siegenian. It has not been found in the reference section or in other spore-bearing strata of the Rhineland (Riegel and Karathanasopoulos, 1982), or in the U.S.S.R.

In the Eifel region, the *annulatus-sextantii* Zone can be subdivided in the Lower Emsian by the appearance of *Verruciretusispora dubia* and possibly *Calyptosporites biornatus* in the lower Klerf Formation. However, *V. dubia* has not been recorded below the succeeding *douglastownense-eurypterota* Zone in Gaspé. *Acinosporites lindlarensis*, a characteristic species of the *douglastownense-eurypterota*

*There is a disagreement between the authors regarding the generic names of some pseudosaccate spores. For those taxa marked with an asterisk DCM would prefer to use *Grandispora* and JBR would prefer *Calyptosporites* or, for some, *Spinozonotriletes*. No new combinations are proposed. For discussion see Richardson (1960, p. 58; 1965, p. 583), Playford (1971, p. 45), Riegel (1973, p. 97), and McGregor and Camfield (1982, p. 43).

Zone in western Europe, occurs high in the *annulatus-sextantii* Zone in the Stooping River Formation. *Emphanisporites annulatus* and some other species of the *annulatus-sextantii* Zone are present in the Icla Formation of the Tarabuco region, Bolivia (McGregor, 1984). However, *E. annulatus* has not been recorded from Arctic Canada or Spitsbergen.

### 10. **Grandispora douglastownense - Ancyrospora eurypterota* Assemblage Zone

***Age.*** Late Emsian and earliest Eifelian. Approximately *Polygnathus costatus patulus* Zone to lowermost *Polygnathus costatus costatus* Zone inclusive, of the conodont scale.

***Reference section for the base of the zone.*** Uppermost Wetteldorf Formation, Plein Quarry, west side of Remel-Bach-Tal, 3 km east of Murlenbach, Salmerwald Syncline, Eifel region (Riegel, 1975, p. 62, 258; 1982).

***Range of the zone.*** High in the Wetteldorf Formation to the top of the Lauch Formation.

***Characteristic species.***

*Grandispora douglastownense* McGregor, 1973, Pl. 8, figs. 8, 9, 12-14 (Pl. 8, fig. 3).

*Ancyrospora eurypterota* Riegel, 1973, Pl. 17, figs. 1-5 (Pl. 8, figs. 1, 2).

*Acinosporites apiculatus* (Streel) Streel, 1967; Streel, 1964, Pl. 1, fig. 13 (Pl. 8, figs. 4, 5).

*Ancyrospora kedoae* (Riegel) Turnau, 1974; Riegel, 1973, Pl. 15, figs. 1-3 (Pl. 9, fig. 1).

*A. nettersheimensis* Riegel, 1973, Pl. 17, figs. 6-8 (Pl. 8, figs. 7, 8).

*A. loganii* McGregor, 1973, Pl. 9, figs. 6, 7, 10-12 (Pl. 8, fig. 6).

*Apiculiretusispora gaspiensis* McGregor, 1973, Pl. 3, figs. 1-4 (Pl. 9, fig. 2).

**Calyptosporites* sp. nov. Riegel, 1975; 1982, Pl. 1, fig. 4 (? = *Hymenozonotriletes endemicus* Chibrikova, 1959) (Pl. 9, fig. 3).

**Calyptosporites proteus* (Naumova) Allen, 1965, Pl. 103, figs. 10, 11 (Pl. 9, fig. 4).

*Dibolisporites echinaceus* (Eisenack) Richardson, 1965; see previous zone.

*Grandispora eximia* (Allen) McGregor and Camfield, 1982, Pl. 10, figs. 2, 6, 7 (Pl, 9, fig. 5).

*Hystricosporites* cf. *H. corystus* Richardson, 1965; Riegel, 1973, Pl. 12, figs. 3, 4 (Pl. 9, fig. 7).

*H. microancyreus* Riegel, 1973, Pl. 12, figs. 6, 7 (Pl. 10, figs. 1, 2).

**Spinozonotriletes arduinnae* Riegel, 1973, Pl. 16, figs. 1-3 (Pl. 9, fig. 6).

***Description.*** Appearance of bifurcate appendages (*Ancyrospora, Hystricosporites*). Proliferation of large, apiculate and spinose, zonate-pseudosaccate spores (*Calyptosporites-Grandispora*). Disappearance of most reticulate and tripapillate spores and distally sculptured forms of *Emphanisporites*. Marked increase in spore size. Persistence and proliferation of *Dibolisporites echinaceus, Apiculatasporites microconus*, and *Verruciretusispora dubia*.

***Distribution and regional variation.*** Upper Wetteldorf Formation and Heisdorf and Lauch formations of the Eifel region (Riegel, 1973, 1982); uppermost Hierges Formation of the Ardennes (Streel, Demaret-Fairon and Otazo, 1975; Streel et al., 1981); upper Battery Point Formation of Gaspé (McGregor, 1977), Wapske Formation of New Brunswick, McAdam Lake Formation of Nova Scotia, Blue Fiord Formation south of Strathcona Fiord, Ellesmere Island, and unnamed strata in the subsurface of Melville Island and the Grand Banks (McGregor, unpublished) in Canada; Trout Valley Formation of Maine, U.S.A. (Andrews et al., 1977); subsurface of Poland (Turnau, 1974). Possibly in North Africa (Massa and Moreau-Benoit, 1976), the Kuznets Basin and the southwest Urals (Chibrikova, 1977).

Bifurcate spines seem to be less common at this level in asiatic parts of the Soviet Union, but the comparison is of questionable validity owing to uncertain correlation. In the Barrandian region, the strata in which this zone would be expected, that is, the upper part of the Třebotov Limestone and its correlatives, contain only rare, poorly preserved spores (McGregor, unpublished). Spores high in the preceding spore zone have been recovered from the tentaculite upper *N. richteri* Zone of the lower Třebotov Limestone at Praha-Hlubočepy (McGregor, 1979a).

***Remarks.*** If *Calyptosporites* sp. nov. of Riegel (see above) can be equated with *Hymenozonotriletes endemicus*, a link would be established with assemblages from the Ryazhsk Horizon of the central part of the Russian Platform (Arkhangelskaya, 1974) and the *Calceola* Horizon of the South Urals (Chibrikova, 1977, fig. 36).

The upper part of the Ryazhsk on the Russian Platform, including the lower *Periplecotriletes tortus* Zone of Soviet nomenclature (see Figure 1) was correlated with the Heisdorf Formation (i.e. the lower part of the *douglastownense-eurypterota* Zone) of the Eifel by Arkhangelskaya (1983). In contrast, the Ryazhsk and *Calceola* deposits of the South Urals were aligned with a higher part of the Eifel sequence by Chibrikova (1983), possibly with the Lauch and Nohn formations. The Ryazhsk and *Calceola* beds would then correlate at least in part with the *velatus-langli* Zone, a conclusion close to Arkhangelskaya's earlier (1976) opinion on correlation of these beds. From photographs and other information in the paper by Arkhangelskaya (1976) it would seem that the thin, upper part of the Ryazhsk Horizon in the central regions of the Russian Platform, with **Calyptosporites velatus* and *?Densosporites devonicus*, would be no older than the Nohn Formation (*velatus-langli* Zone, early Eifelian) of the Eifel. This is in contrast to Arkhangelskaya's later (1983) correlation of the upper Ryazhsk with the Heisdorf Formation (late Emsian).

### 11. *Calyptosporites velatus* - *Rhabdosporites langii* Assemblage Zone

**Age.** Early Eifelian. *Polygnathus costatus costatus* Zone of the conodont succession.

***Reference section for base of zone.*** Nohn Formation, Lierberg section, north side of Sötenich Syncline, Eifel region (Paulus, 1961, p. 37; Riegel, 1975). New exposures being investigated in the north Eifel region may provide a more suitable reference section (W. Riegel, pers. comm., 1981).

***Range of zone.*** Base to near top of Nohn Formation.

***Characteristic species.***

*Calyptosporites velatus* (Eisenack) Richardson, 1962; Richardson, 1960, Pl. 14, fig. 4 (Pl. 10, fig. 6).

*Rhabdosporites langii* (Eisenack) Richardson, 1960, Pl. 14, figs. 8, 9 (Pl. 10, fig. 5).

*Acinosporites acanthomammillatus* Richardson, 1965, Pl. 91, figs. 1, 2 (Pl. 10, figs. 3, 4).

*A. macrospinosus* Richardson, 1965, Pl. 91, figs. 3-6 (Pl. 10, fig. 9).

*Anapiculatisporites petilus* Richardson, 1965, Pl. 89, fig. 11 (Pl. 10, fig. 7).

*Ancyrospora ancyrea* (Eisenack) Richardson var. *ancyrea*; Richardson, 1962, Pl. 25, figs. 6, 7 (Pl. 11, figs. 1, 2).

*Corystisporites multispinosus* Richardson, 1965, Pl. 89, fig. 10 (Pl. 12, fig. 1).

*Densosporites concinnus* (Owens) McGregor and Camfield, 1982, Pl. 6, figs. 13, 14, 18, 19 (Pl. 12, fig. 2).

*Grandispora longa* Chi and Hills, 1976, Pl. 14, figs. 1, 2 only (Pl. 11, figs. 7, 8).

*Kraeuselisporites acerosus* (Arkhangelskaya) McGregor and Camfield, 1982, Pl. 14, figs. 6, 7, 9, 10 (Pl. 11, figs. 3, 4).

*Kraeuselisporites ollii*? (Chibrikova) McGregor and Camfield, 1982, Pl. 14, figs. 2-5, 8 (Pl. 10, fig. 8).

*Periplecotriletes tortus* Egorova, 1974, Pl. 1, fig. 5 (Pl. 11, figs. 5, 6).

*Perotrilites bifurcatus* Richardson, 1962, Pl. 25, figs. 4, 5 (Pl. 12, fig. 3).

*Retusotriletes distinctus* Richardson, 1965, Pl. 88, figs. 7, 8 (Pl. 12, fig. 5).

*R. rugulatus* Riegel, 1973, Pl. 10, figs. 2-4 (Pl. 12, fig. 4).

**Description.** Proliferation of finely-sculptured zonate-pseudosaccate spores intermediate in size between miospores and megaspores. Beginning of spores that are both zonate and camerate and have a flimsy exoexine (*Perotrilites sensu* Evans, 1970). Diversification of murornate-spinose spores (*Acinosporites*). Persistence of most species from the previous zone.

***Distribution and regional variation.*** Upper Lauch and Nohn formations of the Eifel region (Riegel, 1973, 1975, 1982); Morsovo Horizon, Russian Platform (Arkhangelskaya, 1976); upper Battery Point Formation, Gaspé (McGregor, 1977); McAdam Lake Formation of Nova Scotia (McGregor, unpublished); Cape De Bray Formation and lowest Weatherall Formation, Melville Island, Arctic Canada (McGregor and Camfield, 1982); Strathcona Fiord Formation and lowermost Hecla Bay Formation, Ellesmere Island, Arctic Canada (McGregor, unpublished); Aouinet Ouenine Formation of Libya (Massa and Moreau-Benoit, 1976; Moreau-Benoit, 1980). Both nominal species are important elements of early Eifelian to late Givetian assemblages in southern China and in the ORS region across Arctic and Eastern Canada, Western Europe, North Africa, the Russian Platform and probably Central Asia. *Acinosporites* spp. may have a more restricted geographical distribution, but the spores of this genus are common in the Scottish Old Red Sandstone, parts of Eastern and Arctic Canada, and probably the Russian Platform. *Ancyrospora ancyrea* has not so far been recorded from Soviet Asia but has been reported from southern China (Gao and Hou, 1975).

### 12. *Densosporites devonicus* - **Grandispora naumovii* Assemblage Zone

**Age.** Mid-Eifelian to early Givetian. Approximately upper *Polygnathus costatus costatus* Zone to upper *Polygnathus xylus ensensis* Zone inclusive in the conodont scale (by indirect correlation).

***Reference section for the base of the zone.*** Type section of the Weatherall Formation, 152 m above its base, 16.2 km east of the east arm of Weatherall Bay, northeastern Melville Island, Arctic Canada (McGregor and Camfield, 1982). Accessory reference section, upper Nohn Formation, Lierberg section, north side of the Sötenich Syncline, Eifel region (Paulus, 1961; Riegel, 1973).

***Range of the zone.*** Near the base to near the top of the Weatherall Formation, eastern Melville Island. Uppermost Nohn to "Fleringer" and Dreimühlen(?) formations, Eifel region (the position of the upper part of the zone is uncertain in the Eifel).

***Characteristic species.***

*Densosporites devonicus* Richardson, 1960, Pl. 14, fig. 10 (Pl. 12, fig. 6).

*Grandispora naumovii* (Kedo) McGregor, 1973; Kedo, 1955, Pl. 4, fig. 8 (Pl. 13, fig. 1).

*Acinosporites acanthomammillatus* Richardson; see previous zone.

*A. macrospinosus* Richardson; see previous zone.

*Ancyrospora grandispinosa* Richardson, 1960, Pl. 14, fig. 7 (Pl. 13, fig. 7).

***Cristatisporites orcadensis* Richardson, *1960, Pl. 14, fig. 2* (Pl. 13, fig. 2).

*Densosporites inaequus* (McGregor) McGregor and Camfield, 1982, Pl. 7, figs. 1-6 (Pl. 12, fig. 7).

*Grandispora inculta* Allen, 1965, Pl. 734, figs. 7-9 (Pl. 13, fig. 5).

*G. mammillata* Owens, 1971, Pl. 14, figs. 1-4 (Pl. 12, fig. 8).

*Hystricosporites gravis* Owens, 1971, Pl. 8, figs. 1-3 (Pl. 14, fig. 1).

*H. reflexus* Owens, 1971, Pl. 7, figs. 1-4 (Pl. 14, fig. 7).

*Kraeuselisporites rugosus* (Owens) McGregor and Camfield, 1982, Pl. 14, figs. 11, 12 (Pl. 13, fig. 6).

*Lophotriletes devonicus* (Naumova ex Chibrikova) McGregor and Camfield, 1982, Pl. 15, figs. 5-11 (Pl. 13, fig. 3).

*Perotrilites conatus* Richardson, 1965, Pl. 91, figs. 10, 11 (Pl. 13, fig. 4).

*Retispora archaelepidophyta* (Kedo) McGregor and Camfield, 1982, Pl. 16, figs. 4-7, 10 (Pl. 14, figs. 3, 4).

*Retusotriletes distinctus* Richardson, 1965; see previous zone.

*R. rugulatus* Riegel; see previous zone.

*Rhabdosporites langii* (Eisenack) Richardson; see previous zone.

*Verrucosisporites premnus* Richardson, 1965, Pl. 90, figs. 1, 2 (Pl. 14, figs. 5, 6).

*V. scurrus* (Naumova) McGregor and Camfield, 1982, Pl. 18, figs. 10-17, 22 (Pl. 14, fig. 2).

***Description.*** First appearance and diversification of cingulizonate spores with bifurcate spines (*Densosporites* spp.). First appearance of irregular, large, verrucate sculpture (e.g. *Verrucosisporites premnus*, *V. scurrus*). Beginning of foveoreticulate sculpture on pseudosaccate spores (*Retispora*). Persistence of all characteristic species of the previous zone. No sharp distinction in size can yet be made between miospores and megaspores.

***Distribution and regional variation.*** Weatherall Formation of eastern Melville Island (McGregor and Camfield, 1982), Bird Fiord Formation of Ellesmere, Helena and Devon islands, lower Hecla Bay Formation of southeastern Ellesmere Island, and subsurface of Bathurst and Cameron islands (McGregor, unpublished) in the Canadian Arctic; upper Battery Point Formation and Malbaie Formation of Gaspé (McGregor, 1977) and Williams Island Formation of Northern Ontario (McGregor and Camfield, 1976) in Eastern Canada; Ahrdorf, Junkerberg, Freilingen, "Fleringer" and Dreimühlen(?) formations of the Prüm Syncline, and upper Nohn Formation of the Sötenich Syncline, Eifel region (Riegel, 1975); upper Lower and Upper Caithness Flagstone, and upper Lower and lower Upper Stromness groups, Orcadian Basin, Scotland (Richardson, 1965; Richardson, unpublished); Mosolovo and Chernoyar horizons of the Russian Platform (Arkhangelskaya, 1974). Possibly in the McAras Brook Formation of St. George's Bay, Nova Scotia, Canada (McGregor, unpublished).

*Densosporites devonicus* has been found in Scotland, Germany, Arctic and Eastern Canada and central parts of the Russian Platform. In the Rhineland, it is rare and sporadic in the Nohn, Ahrdorf, Junkerberg and Freilingen formations. It is abundant at some localities in the Achanarras Fish Bed, Scotland (Richardson, 1965). In western Europe, *Verrucosisporites premnus* is restricted to the upper part of the zone, and in the Orcadian Basin it has not been recorded below the Eday Group, northeast Scotland (Richardson, 1965).

### 13. *Geminospora lemurata* - *Cymbosporites magnificus* Assemblage Zone

***Age.*** Mid-Givetian. Approximately equivalent to the upper *Polygnathus xylus ensensis* Zone to the lower Upper *Polygnathus varcus* Subzone inclusive, of the conodont scale (by indirect correlation).

***Reference section for the base of the zone.*** Uppermost beds of the Weatherall Formation, northeast of Beverley Inlet, southeastern Melville Island, Arctic Canada (Embry and Klovan, 1976; McGregor and Uyeno, 1972; McGregor, 1981).

***Range of the zone.*** Upper Weatherall Formation, from 31.5 m below its contact with the Hecla Bay Formation, to the upper beds of the Hecla Bay Formation.

***Characteristic species.***

*Geminospora lemurata* Balme, 1962, emend. Playford, 1983, figs. 1-9 (Pl. 4, figs. 8, 9).

*Cymbosporites magnificus* (McGregor) McGregor and Camfield, 1982, Pl. 6, figs. 4, 5; McGregor, 1960, Pl. 13, figs. 2-4 (Pl. 14, fig. 11).

*Ancyrospora langii* (Taugourdeau-Lantz) Allen, 1965, Pl. 106, figs. 5-7 (Pl. 16, figs. 2, 3).

*Aneurospora goensis* Streel, 1964, Pl. 1, figs. 16-20 (Pl. 16, fig. 4).

*A. greggsii* (McGregor) Streel, in Becker et al., 1974; McGregor, 1964, Pl. 1, figs. 1-12 (Pl. 15, fig. 10).

*Archaeozonotriletes variabilis* Naumova, 1953; Allen, 1965, Pl. 100, figs. 3-6 (Pl. 14, fig. 10).

*A. timanicus* Naumova, 1953, Pl. 12, fig. 14 (Pl. 15, figs. 4, 5).

*Biharisporites parviornatus* Richardson, 1965, Pl. 90, figs. 12, 13 (Pl. 15, fig. 1).

*Corystisporites serratus* (Naumova) McGregor and Camfield, 1982; Naumova, 1953, Pl. 1, figs. 19, 20 (Pl. 15, fig. 6).

**JBR prefers to use the name *Samarisporites*.

*Geminospora*? *bislimbata* (Chibrikova) McGregor and Camfield, 1982, Pl. 8, figs. 12, 13; Chibrikova, 1959, *Pl. 8, fig. 8 (Pl. 15, fig. 2).*

*Grandispora tomentosa* Taugourdeau-Lantz, 1967, Pl. 1, figs. 7, 8 (Pl. 16, fig. 8).

*Hymenozonotriletes celeber* Chibrikova, 1959, Pl. 13, fig. 3 (Pl. 15, fig. 7).

*Perotrilites conatus* Richardson; see previous zone.

*P. heclaensis* McGregor and Camfield, 1982, Pl. 15, figs. 18-21 (Pl. 15, fig. 3).

***Description.*** Appearance of small-sculptured, camerate spores with a thick exoexine (*Geminospora*). Proliferation of spores with highly irregular verrucate to baculate sculpture (*Verrucosisporites* spp. and *Dibolisporites* spp.). Diversification of species of patinate spores (*Archaeozonotriletes* spp., *Cymbosporites* spp.). Clear distinction in size between miospores and megaspores.

***Distribution and regional variation.*** Uppermost Weatherall Formation and Hecla Bay Formation of Melville Island (McGregor and Uyeno, 1972; McGregor, 1981) and correlative strata of Bathurst, Vanier, Cameron, and Ellesmere islands (McGregor, unpublished) in the Canadian Arctic; Vorobyev, Staryi Oskol and Ardatov beds and their correlatives of the European U.S.S.R. (Naumova, 1953; Kedo, 1955; Araslanova, 1973; Raskatova, 1974). Possibly in the upper Mimer Valley Formation of Vestspitsbergen (Allen, 1965, 1967) and the Blacourt Formation of the Boulonnais, France (Loboziak and Streel, 1980).

***Remarks.*** *Archaeozonotriletes basilaris* Naumova (1953), regarded by Soviet palynologists as an indicator of Vorobyev (early Givetian) and younger age, is thought by Medianik (1982) to be a junior synonym of *Geminospora lemurata*. *A. basilaris* reaches its acme in strata of Mullino to Kynov age in the U.S.S.R. (Araslanova, 1973; Medianik, 1982), (mid to late Givetian in terms of the stage definitions used in this paper), and continues up to the late Frasnian in lesser numbers. Its range and acme thus are similar to those known on a world scale for *G. lemurata*. *A. basilaris* may also be at least a partial synonym of *Aneurospora greggsii* (McGregor) Streel, but this is yet to be determined.

### 14. *Contagisporites optivus* var. *optivus* – ***Cristatisporites triangulatus* Assemblage Zone

***Age.*** Late Givetian and early Frasnian. Approximately Upper *Polygnathus varcus* Subzone to Lower *Polygnathus asymmetricus* Zone inclusive, in the conodont scale (by indirect correlation).

***Provisional reference section for the base of the zone.*** Hecla Bay Formation, northeast of Beverley Inlet, southeastern Melville Island, Arctic Canada (Embry and Klovan, 1976; McGregor and Uyeno, 1972; McGregor, 1981). Spore-bearing beds are sparse at the base of the zone in this section. A stratigraphic sequence on the east side of Kellett Strait, Melville Island, now being studied by DCM, may be a more suitable reference section.

***Range of the zone.*** Hecla Bay Formation, 872 m above its base, to the Beverley Inlet Formation, 303 m above its base.

***Characteristic species.***

*Contagisporites optivus* (Chibrikova) Owens var. *optivus*; Owens, 1971, Pl. 16, figs. 1-3 (Pl. 16, fig. 1).

*Cristatisporites triangulatus* (Allen) McGregor and Camfield, 1982; Allen, 1965, Pl. 99, figs. 1-6 (Pl. 15, fig. 9).

*Ancyrospora involucra* Owens, 1971, Pl. 24, figs. 5, 6, Pl. 25, figs. 1, 2 (Pl. 15, fig. 8).

*A. langii* (Taugourdeau-Lantz) Allen; see previous zone.

*A. melvillensis* Owens, 1971, Pl. 23, figs. 5-6 (Pl. 16, figs. 6, 7).

*Aneurospora goensis* Streel; see previous zone.

*A. greggsii* (McGregor) Streel; see previous zone.

*Archaeoperisaccus timanicus* Pashkevich, 1964, Pl. 14, figs. 1-4 (Pl. 16, fig. 9).

*Chelinospora concinna* Allen, 1965, Pl. 101, figs. 12-20 (Pl. 16, fig. 10).

*C. ligurata* Allen, 1965, Pl. 102, figs. 1-7 (Pl. 16, fig. 5).

*Geminospora lemurata* Balme emend. Playford, 1983; see previous zone.

*Grandispora tomentosa* Taugourdeau-Lantz; see previous zone.

***Description.*** First appearance of forms with prominent development of zona in the radial regions (*Cristatisporites triangulatus*). Diversification of small-sculptured camerate spores with a thick exoexine (*Geminospora*) and their dominance in some nearshore marine and external continental facies. Waning of *Rhabdosporites langii*, *Calyptosporites velatus* and *Dibolisporites echinaceus*. Disappearance of *Grandispora naumovii* and *Perotrilites bifurcatus*. Persistence, and abundance in some assemblages, of anchor-spined spores with broad, conical spines that bear small, terminal bifurcations (e.g. *Ancyrospora melvillensis*). Appearance of multifurcate spines high in the zone (Taugourdeau-Lantz, 1971). First monolete camerate spores (*Archaeoperisaccus timanicus*) in the middle of the zone (Medianik, 1980; McGregor, 1981).

***Distribution and regional variation.*** Hecla Bay and Beverley Inlet formations of eastern Melville Island, Weatherall Formation of western Melville Island and Prince Patrick Island (McGregor, 1981), upper Hecla Bay Formation and lower Fram Formation of southwestern Ellesmere Island, Hare Indian Formation of the western District of Mackenzie, lower Flume Formation of east central British Columbia, Escuminac Formation of Quebec, and unnamed strata in the subsurface of the Grand Banks (McGregor, unpublished) in Canada; subsurface of Belgium (Streel, 1972); Blacourt Formation, Boulonnais, France (Loboziak and Streel, 1980); upper Mimer Valley Formation of Vestspitsbergen (Allen, 1965, 1967); and the Mullino, Pashiya, Kynov and Sargaevo beds and their correlatives in the European U.S.S.R. (Chibrikova, 1962; Araslanova, 1973). Possibly in the Nation River Formation of east-central Alaska (Scott and Doher,

1967), the Pine Point Formation of the District of Mackenzie, Canada (Audretsch, 1968), the subsurface of southern England (Mortimer and Chaloner, 1972; McGregor, 1981), the Sherkin Formation of County Cork, Ireland (Clayton and Graham, 1974), and the Bu Ness, Observatory and Ward Hill groups of Fair Isle, Shetland (Marshall and Allen, 1982).

The "Haikou Formation" of Longhuashan, Zhanyi, China contains *Cristatisporites* (al. *Samarisporites*) *triangulatus* and *Chelinospora ligurata*, and a *Stringocephalus* fauna that is taken to indicate Givetian age (Lu, 1980a). It also contains a distinctive monolete spore superficially like *Archaeoperisaccus* but with transverse exinal folds suggesting distal cavation (Lu, 1980b). Spores like it have not been found in the ORS region. Two other species in the same spore assemblage (Lu, 1980b) apparently are similar to *Archaeoperisaccus* spp. of the ORS region and may represent the earliest record of specimens of this genus.

***Remarks.*** The appearance of multifurcate spines and *A. timanicus* may eventually form the basis for subdividing this zone.

Loboziak and Streel (1980) identified *Cristatisporites triangulatus* from near the base of the Blacourt Formation in the Boulonnais, immediately below conodonts of the *Icriodus obliquimarginatus* Zone. As far as we are aware, this is the earliest age claimed for this species. The age of the sample seems to be beyond dispute. If their concept of *C. triangulatus* is the same as ours, and/or as that of the Soviet palynologists (see Pychova, 1960; Volochaev et al., 1979; Kedo and Obukhovskaya, 1981), the documented stratigraphic range of this species would be extended significantly downward by this record, and the base of the *optivus-triangulatus* Zone would need to be redefined.

However, the oldest specimens identified as *C. triangulatus* in the Boulonnais occur three conodont zones below the first record of *Geminospora lemurata* (Loboziak and Streel, 1980) in the same region, whereas in Arctic Canada (McGregor, 1981), and in the European U.S.S.R. (e.g. in Fomina, 1968), their order of appearance is reversed, if one uses the synonymy for *G. lemurata* given in McGregor and Camfield (1982) and suggested by Playford (1983), and the synonymy for *C. triangulatus* given by Allen (1982). The context of the assemblages in which these species first occur in Canada and the U.S.S.R. suggests that *G. lemurata* may occur well below *C. triangulatus* in beds equivalent to the conodont *obliquimarginatus* Zone (Fig. 6). Nevertheless, none of the Canadian or Soviet reports include data on co-occurring faunas that would permit international correlation of the spore ranges. Further work is needed in order to correlate the stratigraphic ranges of both species.

### 15. *Archaeoperisaccus ovalis - Verrucosisporites bulliferus* Assemblage Zone

***Age.*** Frasnian. Approximately the Middle *Polygnathus asymmetricus* Zone to the *Palmatolepis gigas* Zone inclusive in the conodont scale.

***Reference section for the base of the zone.*** Noces Member, Beaulieu Formation, Caffiers area, Ferques railroad section, Boulonnais, France (Brice et al., 1979; Loboziak and Streel, 1981). The base of this zone is well defined in this section, as it is in stratigraphical continuity with beds below containing spores of the *optivus-triangulatus* Zone. Accessory reference section (for the *Archaeoperisaccus* assemblage), Beverley Inlet Formation, northeast of Beverley Inlet, southeastern Melville Island (Embry and Klovan, 1976; McGregor and Uyeno, 1972; McGregor, 1981).

***Range of the zone.*** In the Boulonnais, from the Noces Member of the Beaulieu Formation to the upper (but not uppermost) Hydrequent Formation inclusive. In the Canadian Arctic, from the Beverley Inlet Formation, 353 m above the base, to about 110 m above the base of the overlying Parry Islands Formation.

***Characteristic species.***

*Archaeoperisaccus ovalis* Naumova, 1953, Pl. 14, fig. 13 (Pl. 17, fig. 1).

*Verrucosisporites bulliferus* n. name; replaced synonym, *V. bullatus* Taugourdeau-Lantz, 1967, p. 50, Pl. 2, figs. 9, 10 (non *V. bullatus* Balme and Hennelly, 1956) (Pl. 17, fig. 2).

*Ancyrospora furcula* Owens, 1971, Pl. 23, figs. 1-4 (Pl. 17, figs. 4, 5).

*A. melvillensis* Owens, 1971; see previous zone.

*Aneurospora greggsii* (McGregor) Streel, 1974; see previous two zones.

*Archaeoperisaccus opiparus* Owens, 1971, Pl. 22, figs. 1-6 (Pl. 17, fig. 3).

cf. *Archaeozonotriletes laetus* Sergeeva in Nazarenko et al., 1971, Pl. 5, fig. 11 (Pl. 17, fig. 7).

*Diaphanospora reticulata* Guennel, 1963, figs. 17-19 (Pl. 17, fig. 11).

*Geminospora lemurata* Balme emend. Playford, 1983; see previous two zones.

*Hymenozonotriletes deliquescens* Naumova, 1953, Pl. 9, fig. 8 (Pl. 17, fig. 6).

*Lophozonotriletes media* Taugourdeau-Lantz, 1967, Pl. 2, fig. 6 (Pl. 17, fig. 10).

*Nikitinsporites canadensis* Chaloner, 1959, Fig. 4, Pl. 55, fig. 5 (Pl. 17, figs. 8, 9).

***Description.*** First appearance and acme of minutely sculptured, monolete, camerate spores (*Archaeoperisaccus* spp). Proliferation of multifurcate-spined spores (e.g. *Ancyrospora furcula*) and small megaspores (e.g. *Nikitinsporites*). Most ancyrospores with broad, conical, bifurcate spines disappear in the upper part of the zone. Continued prominence of *Geminospora* spp. and bifurcate-spined spores. Decline and disappearance of most large, sculptured, zonate-pseudosaccate miospores. Beginning of marked regionalism in the distribution of some elements of the spore floras.

***Distribution and regional variation.*** Beaulieu, Ferques and Hydrequent formations, Boulonnais, France (Loboziak and Streel, 1981); Beverley Inlet Formation and lower Parry




***Figure 6.*** *Approximate correlation of selected late Givetian and early Frasnian spore-bearing strata of the European U.S.S.R., Western Europe, and Arctic Canada.*

Islands Formation, southeastern Melville Island (Owens, 1971; McGregor and Uyeno, 1972; McGregor, 1981), subsurface of Vanier Island (Whiteley, 1980), Okse Bay Group of Grinnell Peninsula, Devon Island, and subsurface of Banks and Cameron islands (McGregor, unpublished), Fram and Hell Gate formations of southwestern Ellesmere Island (McGregor, 1969 and unpublished), Weatherall Formation of Prince Patrick Island (McGregor, 1981), Imperial Formation of District of MacKenzie (McGregor, 1969; Chi and Hills, 1974), Escarpment Member of Hay River Formation, District of MacKenzie (McGregor and Owens, 1966) and Perdrix Formation of east-central British Columbia (McGregor, unpublished) in Canada; Domanik, Liaiol, Vetlasyan, Bel'gop and Sirachoi strata of the Timan-Pechora region (Sennova, 1972; Medianik, 1980), Goltva and Rudnya strata of the Dnepr-Donets Depression (Sergeeva, 1979), and Semiluki, Petino, Voronezh and Evlanov strata and their correlatives on the Russian Platform (Naumova, 1953; Raskatova, 1974) in the U.S.S.R.; Chemung Group of central Pennsylvania, U.S.A. (McGregor, unpublished). Possibly in the Calmar Formation of western Alberta and the Long Rapids Formation of Northern Ontario, Canada (McGregor, unpublished); the Oneonta, Walton and Rhinestreet formations of central New York State, U.S.A. (Richardson, unpublished); and the subsurface of the Vilyu River region in east-central Siberia (Mikhailova and Fradkina, 1966).

The nominal species ***Verrucosisporites bulliferus***, present in Northern Ontario (McGregor, unpublished), central New York State (Richardson, unpublished) and France (Taugourdeau-Lantz, 1971) may be identical to ***Lophozonotriletes grumosus*** Naumova (1953), which is common in middle and late Frasnian deposits of the Russian Platform.




***Figure 7.*** *Approximate correlation of the* ***torquata-gracilis/flexuosa-cornuta*** *boundary in New York State, based on spores and invertebrate faunas. A = ammonoids, C = conodonts, after Rickard (1975) and Kirchgasser and House (1981). The Belgian units Fa2a to Fa2c are diachronous (Becker et al., 1974). We have shown the oldest age for each as indicated by Ziegler (1979).*

*Archaeoperisaccus* spp. apparently are geographically restricted, as they occur in the eastern, northern and western parts of the ORS region and in eastern Asia (McGregor, 1979b, Fig. 13).

### 16. *Auroraspora torquata* - *Grandispora gracilis* Assemblage Zone

**Age.** Latest Frasnian to early late Famennian. Approximately the lower *Palmatolepis triangularis* Zone to the middle *Scaphignathus velifer* Zone inclusive in terms of the conodont succession (Rickard, 1975; for post-Gowanda formations of New York, unpublished spore correlation by JBR with the Belgian succession). See Figure 7 and comments under the *flexuosa-cornuta* Zone.

***Reference section for the base of the zone.*** Hanover Shale Formation, West Falls Group, Chautauqua County, western New York State, U.S.A., Walnut Creek section (Pepper and de Witt, 1950; Ahmed, 1978; Richardson, unpublished).

***Range of the zone.*** Hanover Shale Formation, 9.8 m above its base, to the lower part of the upper Northeast Shale Formation, inclusive.

***Characteristic species.***

*Auroraspora torquata* Higgs, 1975, Pl. 4, figs. 1-3 (Pl. 18, fig. 1).

*Grandispora gracilis* (Kedo) Streel, in Becker et al., 1974, Pl. 19, figs. 1-3 (Pl. 18, figs. 2, 3).

*Aneurospora greggii* (McGregor) Streel, 1974; see previous three zones.

*Archaeozonotriletes famenensis* Naumova, 1953, Pl. 17, figs. 31-34 (Pl. 18, fig. 4).

*Auroraspora hyalina* (Naumova) Streel, in Becker et al., 1974; Streel, 1965, Pl. 1, figs. 7, 8 (Pl. 18, fig. 6).

*A. macra* Sullivan, 1968, Pl. 27, figs. 6-10 (Pl. 19, fig. 8).

*A. pseudocrista* Ahmed, 1980, Pl. 2, figs. 5-9 (Pl. 18, fig. 8).

*Cornispora varicornata* Staplin and Jansonius in Staplin, 1961, Pl. 1, figs. 6-12 (Pl. 18, fig. 5).

*Crassispora catenata* Higgs, 1975, Pl. 2, figs. 1-3 (Pl. 19, fig. 5).

*Cyrtospora cristifera* (Luber) Van der Zwan, 1979, Pl. 1, figs. 1-5; Pl. 2, figs. 1-8; Pl. 3, figs. 1-3 (Pl. 18, fig. 7).

*Diducites poljessicus* (Kedo) Van Veen, 1981, Pl. 4, figs. 1-4, 6 (Pl. 19, fig. 4).

*D. versabilis* (Kedo) Van Veen, 1981, Pl. 2, figs. 5, 6; Pl. 3, figs. 1-6, 9 (Pl. 18, fig. 9).

"(?)*Hymenozonotriletes denticulatus* Naumova" of McGregor, 1967, Pl. 1, fig. 25 (Pl. 19, figs. 2, 3).

*Hystricosporites multifurcatus* (Winslow) Mortimer and Chaloner, 1967; Winslow, 1962, Pl. 12, figs. 8-8b, Pl. 13, figs. 1-9, Pl. 22, fig. 16 (Pl. 18, fig. 10).

*Knoxisporites dedaleus* (Naumova) Streel, 1977; Naumova, 1953, Pl. 19, fig. 11. [=*K. literatus* (Waltz) Playford, of Clayton et al., 1978)] (Pl. 19, fig. 1).

*Lophozonotriletes lebedianensis* Naumova, 1953, Pl. 17, fig. 42 (Pl. 19, fig. 6).

**Description.** Diversification of smooth-walled, pseudosaccate spores (*Auroraspora*) and abundance of *Aneurospora greggsii*. First appearance of broad radial and/or concentric thickenings on cingulate spores (*Knoxisporites*). First appearance of prominent spinose to verrucate distal crassitude (*Cyrtospora cristifera*). Specimens with two or more protrusions on the crassitude (*Cornispora varicornata*) appear about midway in the zone in northern and western regions. Miospores on average smaller than in Middle Devonian and earlier Frasnian zones, commonly less than 100 µm in diameter.

**Distribution and regional variation.** Parry Islands Formation of Banks, Cameron and Byam Martin islands (McGregor, unpublished) and Bathurst Island (McGregor and Owens, 1966), Kotcho and Imperial formations of western District of Mackenzie, and Memramcook Formation of southeastern New Brunswick (McGregor, unpublished) in Canada; Hanover, Dunkirk, South Wales, Gowanda, Laona, Westfield and lower Northeast Shale formations of western New York State (Ahmed, 1978; Wood, 1978; Richardson, unpublished); Livny, Zadonsk and Elets strata of the Russian Platform (Raskatova, 1973, 1974; Naumova, 1953); Hydrequent Formation, Boulonnais, France (Loboziak and Streel, 1981; Loboziak et al., 1983).

*Auroraspora torquata* is present in Southern Ontario, New Brunswick and Nova Scotia in Canada (McGregor, unpublished) and in New York State and western Europe, but has not been reported from the Soviet Union or the northern parts of the ORS continent. *Cyrtospora cristifera* does not appear in the southern part of the ORS continent until the *nitidus-vallatus* Zone (latest Devonian).

**Remarks.** *Hystricosporites multifurcatus* as used herein does not include the specimens described under this name by Mortimer and Chaloner (1967). Nevertheless, the nomenclatural transfer of the species to *Hystricosporites* as proposed by them is retained.

### 17. *Rugospora flexuosa - Grandispora cornuta* Assemblage Zone

**Age.** Late Famennian. Approximately Upper *Scaphignathus velifer* Zone to lower part of Middle *Bispathodus costatus* Zone inclusive, in terms of the conodont succession and based on spore evidence. This dating does not agree with the age determined on the basis of sparse conodont and ammonoid evidence in New York State (see below and Figure 7).

Rickard (1975, p. 10) records that conodont species of the *Palmatolepis marginifera* Zone have been recognised in the Northeast Shale Formation and overlying "Chadakoin" (Dexterville and Ellicott formations of the Conneaut Group). Kirchgasser and House (1981) record "fauna 27" in the middle of the Ellicott Shale Formation, including the ammonoid *Maeneceras* cf. *pompeckji*, which in Europe is a "typical Famennian II beta goniatite". According to Ziegler (1979), the upper part of the *Cheiloceras* Stufe (II beta) is equivalent to the *Palmatolepis rhomboidea* and lower part of the *P. marginifera* conodont zones. Thus the conodont and ammonoid evidence, although scanty, is more or less consistent.

In contrast, a correlation (miospore-based) with Belgian strata where conodont sequences are well known, would place the appearance of *Rugospora flexuosa* and *Grandispora cornuta* (as *G. uncata*) at or near the base of Fa2c (Becker et al., 1974), a level which, according to Ziegler (1979, Fig. 7), is not older than the base of the conodont Upper *S. velifer* Zone. In Belgium, the equivalent of the upper part of the *flexuosa-cornuta* Zone (the upper VCo Zone of Streel) has been correlated with the lower part of the conodont Middle *B. costatus* Zone (see enclosure in Paproth et al., 1983). Therefore, on spore evidence, the *flexuosa-cornuta* Zone in Belgium may extend from the Upper *S. velifer* Zone to the Middle *B. costatus* Zone of the conodont succession.

It seems unlikely that the discrepancy in New York between the conodont - ammonoid and miospore correlations is due to the late appearance or incomplete record of the nominal miospore species in Belgium. This conclusion is supported by data on the distribution of two other species that have restricted stratigraphic ranges in both Belgium and New York State. In Belgium, *Retusotriletes phillipsii* is confined to lower Fa2c and *Lagenicula* sp. A is only recorded from Fa2c (Becker et al., 1974). In New York State and northern Pennsylvania, if the conodont - ammonoid evidence were accepted, the ranges of these species would be approximately lower and middle Fa2b (*R. phillipsii*) and Fa2a(?) and lower Fa2b (*Lagenicula* sp. A), not Fa2c.

Clendening et al. (1980) also suggest a correlation between the New York State succession and that of Belgium. On the basis of the appearance of *Retusotriletes phillipsii*, they correlate the Shumla Siltstone and Northeast Shale formations of the upper part of the Canadaway Group (below the *flexuosa-cornuta* Zone) with the Belgian Fa2c. This correlation would lower the base of Fa2c even more, in terms of the New York succession, than would the palynological

correlation discussed in the preceding paragraphs. On the other hand, in samples examined by JBR through the western New York State sequence, *R. phillipsii* has not been found below the Dexterville Siltstone Formation, which overlies the Northeast Shale Formation. *Grandispora cornuta* occurs commonly in the Dexterville Formation and above, and a similar form occurs sporadically in the upper part of the underlying Northeast Shale Formation.

***Reference section for the base of the zone.*** North Chautauqua County, New York State, tributary of Canadaway Creek, approximately 2.8 km southeast of Arkwright Falls; upper Northeast Shale Formation (Canadaway Group), 16.8 m above the base of the exposed section, 125 m above the top of the Shumla Siltstone Formation and about 25.6 m below the base of the Dexterville Formation. The exposure is marked on the geological map in Tesmer (1963).

In sections at Chautauqua Creek to the southwest, assemblages belonging to this zone occur in at least the top 17 m of the Northeast Shale Formation (Richardson, unpublished).

No suitable reference section has been found yet for the *Cornispora* assemblage.

***Range of the zone.*** Upper Northeast Shale Formation (Canadaway Group) to the lower Venango and Cattaraugus Formations, western New York State.

***Characteristic species.***

*Rugospora flexuosa* (Jushko) Streel in Becker et al., 1974, Pl. 21, figs. 8-11 (Pl. 19, fig. 7).

*Grandispora cornuta* Higgs, 1975, Pl. 4, figs. 4-6 (Pl. 20, fig. 1).

*Auroraspora hyalina* (Naumova) Streel; see previous zone.

*A. macra* Sullivan; see previous zone.

*A. torquata* Higgs; see previous zone.

*Cornispora varicornata* Staplin and Jansonius; see previous zone.

*Cyrtospora cristifera* (Luber) Van der Zwan; see previous zone.

*Diducites poljessicus* (Kedo) Van Veen; see previous zone.

*D. versabilis* (Kedo) Van Veen; see previous zone.

*Grandispora gracilis* (Kedo) Streel; see previous zone.

*Hystricosporites multifurcatus* (Winslow) Mortimer and Chaloner, 1967; see previous zone.

*Lagenicula* sp. A Streel, in Becker et al., 1974, Pl. 23, figs. 3-7 (=*Hymenozonotriletes immensus* Nazarenko and Nekryata, in Kedo et al., 1971, Pl. 16, figs. 1-4) (Pl. 19, fig. 9).

*Retispora cassicula* (Higgs) Higgs and Russell, 1981; Higgs, 1975, Pl. 5, figs. 1-3 (Pl. 20, fig. 2).

*Retusotriletes phillipsii* Clendening, Eames and Wood, 1980, Pl. 1, figs. 1-9 (Pl. 20, fig. 3).

***Description.*** Reappearance of pseudosaccate spores with foveoreticulate exoexine (*Retispora*; see Zone 12). Continuation of most species from the previous zone, some in greater abundance (e.g. *Auroraspora hyalina, A. macra, A. poljessica, A. torquata, Grandispora gracilis, Cyrtospora cristifera,* and *Rugospora versabilis*). Diversification of simple-spined and laevigate pseudosaccates (*Grandispora* and *Auroraspora*).

***Distribution and regional variation.*** Northeast Shale, Dexterville and Ellicott Formations (upper Canadaway and Conneaut Groups) and lower Venango and Cattaraugus Formations (lower Conewango Group), New York State (Ahmed, 1978; Richardson, unpublished), and an unspecified stratigraphic unit in West Virginia (McGregor, unpublished) in the U.S.A.; Imperial Formation of the Yukon Territory and western District of Mackenzie (Van der Zwan and Walton, 1981; McGregor, unpublished) and Parry Islands Formation of Bathurst and Banks islands, and unnamed strata of Antigonish County, Nova Scotia (McGregor, unpublished) in Canada; Misery Series of Bear Island (Kaiser, 1971); Assise d'Evieux, Belgium (Becker et al., 1974); Purple Sandstone Group of southern Ireland (Higgs and Russell, 1981); subsurface of North Africa (Massa and Moreau-Benoit, 1976); subsurface of Poland (Turnau, 1979); Dankov-Lebedyan beds of the Russian Platform (Raskatova, 1973, 1974; Byvsheva, 1980), the Pripyat Depression (e.g. Kedo et al., 1971; Avkhimovich, 1974), the Dniepr-Donets Depression (Sorokina, 1968) and northern Timan (Pashkevich, 1971; Sennova, 1972) in the U.S.S.R. *Cornispora* spp. and *Cyrtospora cristifera* are restricted to the eastern, northern and western parts of the ORS Continent, as in the previous zone.

### 18. *Vallatisporites pusillites* (sensu lato) – *Retispora lepidophyta* Zone

***Age.*** Strunian (Conil and Lys, 1980). Approximately Middle *Bispathodus costatus* Zone to lower part of the "Lower *Protognathodus* fauna" in terms of the German conodont succession.

***Reference section for the base of the zone.*** Chanxhe Section, eastern Dinant Basin, Ourthe Valley, Belgium (Bless et al., 1974; Streel, Bless et al., 1975).

***Range of the zone.*** Fa2d to lower Tn1b. For more extensive discussion of the range of this zone, see Clayton et al. (1977, p. 6).

***Characteristic species.***

*Vallatisporites pusillites* (Kedo) Dolby and Neves, 1970, Pl. 2, figs. 1-4 (Pl. 20, fig. 5).

*Retispora lepidophyta* (Kedo) Playford, 1976, Pl. 10, figs. 1-15 (Pl. 20, fig. 4).

*Diducites poljessicus* (Kedo) Van Veen; see previous two zones.

*D. versabilis* (Kedo) Van Veen; see previous two zones.

*Discernisporites micromanifestus* (Hacquebard) Sabry and Neves, sensu Higgs, 1975, Pl. 5, fig. 13 (Pl. 20, fig. 7).

*Grandispora echinata* Hacquebard, 1957, Pl. 3, fig. 17 (Pl. 20, fig. 6).

*Knoxisporites literatus* (Waltz) Playford, 1963, Pl. 90, figs. 7, 8 (Pl. 21, fig. 5).

*Raistrickia variabilis* Dolby and Neves, 1970, Pl. 1, fig. 6 (Pl. 21, fig. 1).

*Rugospora flexuosa* (Jushko) Streel; see previous zone.

*Tumulispora malevkensis* (Kedo) Turnau, 1978; Kedo, 1963, Pl. 10, figs. 240, 241 (Pl. 21, fig. 4).

*T. rarituberculata* (Luber) Turnau, 1975, Pl. 4, figs. 1, 2, 4, 6 (Pl. 21, figs. 2, 3).

**Description.** Acme of pseudosaccate spores with a foveoreticulate exoexine (*Retispora*). Widespread occurrence of vacuolate zona (*Vallatisporites*). Persistence of most of the species characteristic of the previous zone.

**Distribution and regional variation.** This zone is of near worldwide occurrence (see McGregor, 1979b). In the standard North American succession in western New York State, *Vallatisporites pusillites* begins in the upper part of the preceding *flexuosa-cornuta* Zone, in the lowermost parts of the Venango and Cattaraugus formations about 79 m below the appearance of *Retispora lepidophyta*. It also begins in the upper part of the *flexuosa-cornuta* Zone in the Lower Sandstone Member of the Pocono Formation of Pennsylvania (Streel and Traverse, 1978). In Belgium, *V. pusillites* and *R. lepidophyta* make their first appearance together, at the base of the *pusillites-lepidophyta* Zone. The two species also occur together in Kazakhstan (Grishina and Klenina, 1981), Tibet (Gao, 1983) and Kwangsi, South China (Wang et al., 1978), but *V. pusillites* is absent from synchronous strata in Hunan (Hou, 1982). On the other hand, in the European U.S.S.R. both *V. pusillites* and *Rugospora flexuosa* appear high in the range of *R. lepidophyta*, probably in the *lepidophyta-nitidus* Subzone (Byvsheva, 1980). *Tumulispora malevkensis* first occurs in this zone in eastern Europe (Byvsheva, 1980; Raskatova, 1974; Turnau, 1978) but not until the succeeding *nitidus-vallatus* zone in western Europe (Clayton et al., 1977).

*Hymenozonotriletes explanatus* is first seen near the top of the zone in southern Ireland, and is used by Clayton et al. (1978) to define the boundary between two subzones in that region: the *lepidophyta-literatus* Subzone and the *lepidophyta-explanatus* Subzone. The *lepidophyta-explanatus* Subzone is characterized by the same species as the *lepidophyta-literatus* Subzone, with the addition of *H. explanatus*. It has been recognized in southern Ireland (Clayton, in Naylor, 1975; Higgs, 1975), the Ardennes-Rhine region, and western New York State and northern Pennsylvania (Richardson, unpublished). However, as *H. explanatus* occurs only in the highest beds of the *pusillites-lepidophyta* Zone, it does not seem reasonable to regard it as a characteristic species of this zone.

**Remarks.** It may be possible to separate a new zone in the upper part of the *pusillites-lepidophyta* Zone based on *H. explanatus*, *Grandispora echinata*, *Knoxisporites literatus*, *Tumulispora* spp. and other species that have not been found in the lower part of the zone.

### 19. *Verrucosisporites nitidus* - *Vallatisporites verrucosus* Assemblage Zone

#### *Retispora lepidophyta* - *Verrucosisporites nitidus* Subzone

**Age.** Late Strunian. Approximately equivalent to all but the basal part of the "Lower *Protognathodus* fauna" of the German conodont succession.

**Reference section for the base of the zone.** Old Head of Kinsale, South Munster Basin, southwestern Ireland (Clayton et al., 1974). The base of the zone is about two thirds of the way up the Holeopen Bay Member of the Old Head Sandstone Formation.

**Range of the subzone.** Upper third of the Holeopen Bay Member to the top of the Old Head Sandstone Formation.

**Characteristic species.**

*Verrucosisporites nitidus* (Naumova) Playford, 1964, Pl. 3, figs. 3-6 (Pl. 21, fig. 7).

*Vallatisporites verrucosus* Hacquebard, 1957, Pl. 2, fig. 13 (Pl. 21, fig. 8).

*Retispora lepidophyta* (Kedo) Playford; see previous zone.

*Cyrtospora cristifera* (Luber) Van der Zwan; see *torquata-gracilis* and *flexuosa-cornuta* zones.

*Discernisporites micromanifestus* (Hacquebard) Sabry and Neves, sensu Higgs, 1975; see previous zone.

*Grandispora echinata* Hacquebard, 1957; see previous zone.

*Hymenozonotriletes explanatus* (Luber) Kedo, 1963, Pl. 6, figs. 144-147 (Pl. 21, fig. 6).

*Knoxisporites literatus* (Waltz) Playford; see previous zone.

*Raistrickia corynoges* Sullivan, 1968, Pl. 25, figs. 6-8 (Pl. 21, fig. 9).

*Tumulispora malevkensis* (Kedo) Turnau; see previous zone.

*T. rarituberculata* (Luber) Turnau; see previous zone.

**Description.** Decline of grapnel-tipped spines (*Ancyrospora*, *Hystricosporites*) and foveoreticulate exoexine (*Retispora*), and their disappearance at the top of the zone. Diversification of coarsely verrucate, cingulate spores (*Tumulispora*). Appearance of *Cyrtospora cristifera* in southern parts of the ORS Continent (eastern North America and western Europe).

**Distribution.** Old Head Sandstone Formation, southern Ireland (Clayton et al., 1974; Holland, 1981); highest Hangenberg Shales, Germany (Paproth and Streel, 1970); unnamed strata on St. George's Bay, Nova Scotia, Banff Formation of Alberta and southwestern District of Mackenzie, and Besa River Formation of east central British Columbia in Canada (McGregor, unpublished); Knapp Formation of western New York State and northern

Pennsylvania (Richardson, unpublished) and Pocono Formation of Pennsylvania, U.S.A. (Streel and Traverse, 1978); Malevka Horizon of the Russian Platform (Byvsheva, 1976). Possibly in the subsurface of Poland (Turnau, 1979), the Sappington Member of Three Forks Formation of Montana (Sandberg et al., 1972) and the Berea Formation of Ohio (Winslow, 1962).

## REFERENCES


Ahmed, S.

1978: Palynology and biostratigraphy of some Upper Devonian deposits of western New York State and northern Pennsylvania, U.S.A.; unpublished Ph.D. thesis, King's College, London, 356 p.

1980: Some forms of the genus *Auroraspora* from the Upper Devonian of New York State and Pennsylvania, U.S.A.; Journal of the University of Kuwait (Science), v. 7, p. 227-239.

Aldridge, R.J., Dorning, K.J., Hill, P.J., Richardson, J.B., and Siveter, D.J.

1979: Microfossil distribution in the Silurian of Britain and Ireland; in The Caledonides of the British Isles – reviewed, A.L. Harris, C.H. Holland, and B.E. Leake (eds.); Geological Society of London, Special Publication no. 8, Scottish Academic Press Ltd., p. 433-438.

Allen, J.R.L.

1979: Old Red Sandstone facies in external basins, with particular reference to southern Britain; in The Devonian System, M.R. House, C.T. Scrutton, and M.G. Bassett, (eds.); Palaeontological Association, Special Paper 23, p. 65-80.

Allen, J.R.L. and Dineley, D.L.

1976: The succession of the Lower Old Red Sandstone (Siluro-Devonian) along the Ross-Tewkesbury Spur Motorway (M50), Hereford and Worcester; Geological Journal, v. 11, no. 1, p. 1-14.

Allen, J.R.L. and Tarlo, L.B.

1963: The Downtonian and Dittonian facies of the Welsh Borderland; Geological Magazine, v. 100, no. 2, p. 129-155.

Allen, K.C.

1965: Lower and Middle Devonian spores of north and central Vestspitsbergen; Palaeontology, v. 8, pt. 4, p. 687-748.

1967: Spore assemblages and their stratigraphical application in the Lower and Middle Devonian of north and central Vestspitsbergen; Palaeontology, v. 10, pt. 2, p. 280-297.

Amoco Canada Petroleum Company Ltd.

1975: Well history report, Amoco Imp. Skelly Phalarope P-62, offshore, Grand Banks, Newfoundland; released January 5, 1977 by the Department of Energy, Mines and Resources, Ottawa, Canada.

Anderton, R., Bridges, P.H., Leeder, M.R., and Sellwood, B.W.

1979: A dynamic stratigraphy of the British Isles. A study in crustal evolution; London, George Allen and Unwin, 301 p.

Andreeva, Ye. M.

1973: Early and Middle Devonian spores from the northeastern slope of Salair; in Stratigraphy of the Lower and Middle Devonian; Academy of Sciences of USSR, VSEGEI, Leningrad, v. 2, p. 40-44, 287-288, 294-295 (in Russian).

Andrews, H.N., Kasper, A.E., Forbes, W.H., Gensel, P.G., and Chaloner, W.G.

1977: Early Devonian flora of the Trout Valley Formation of northern Maine; Review of Palaeobotany and Palynology, v. 23, p. 255-285.

Araslanova, R.M.

1973: Stratigraphical distribution of spores in Devonian terrigenous deposits of the Perm Region of Kama; in Geology, Petroleum and Natural Gas Resources of Northern Part of Ural-Volga Region, S.H. Shikhov (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, No. 123, p. 182-190 (in Russian).

Aristova, K.E. and Arkhangelskaya, A.D.

1976: Microfossils from stratotype sections of Upper Silurian horizons of Estonia; in Results of palynological research on Precambrian, Paleozoic and Mesozoic of the USSR, T.V. Byvsheva (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, No. 192, p. 28-38 (in Russian).

Arkhangelskaya, A.D.

1974: Spore-based zonation of the lower part of the Middle Devonian in central and eastern European parts of the USSR; in Palynology of Proterophyte and Paleophyte, Proceedings of III International Palynological Conference, S.N. Naumova (ed.); "Nauka", Moscow, p. 56-59 (in Russian).

1976: Basis for Eifelian age of the *Periplecotriletes tortus* zone in central districts of the European part of the USSR; in Results of palynological research on Precambrian, Paleozoic and Mesozoic of the USSR, T.V. Byvsheva (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, No. 192, p. 39-66 (in Russian).

1978: Spores from the Lower Devonian of the Lithuanian SSR; Paleontological Journal, v. 1978, no. 2, p. 113-120 (in Russian).

1980a: Plant spores from some Lower Devonian sections of the western regions of the Russian Plate; in Palynological research in the Proterozoic and Phanerozoic of oil and gas bearing regions of USSR, T.V. Byvsheva (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, no. 217, p. 26-46. (in Russian).

1980b: Establishment of the *Retusotriletes clandestinus* Zone in the lower part of the Devonian of the southeastern district of Orenburg Province; in Palynological research in the Proterozoic and Phanerozoic of oil and gas bearing regions of the USSR, T.V. Byvsheva (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, no. 217, p. 47-52 (in Russian).

Arkhangelskaya, A.D. (cont.)
1983: Adjustment of the boundaries of series and stages, and interregional correlation of the lower part of the Devonian of the Russian Platform by horizons; in The Middle Devonian of the USSR, its boundaries and differentiation into stages, M.A. Kamaletdinov (ed.); Academy of Sciences of the USSR, Bashkirian Branch, Institute of Geology, Ufa, p. 74-75 (in Russian.)

Audretsch, A.P.
1968: Middle Devonian microflora from the Great Slave Lake area, Northwest Territories, Canada; in International Symposium on the Devonian System, Calgary, 1967, D.H. Oswald (ed.); Proceedings, v. 2, p. 837-847.

Avkhimovich, V.I.
1974: Palynological characterization of the upper Famennian saliferous formation in the Pripyat Depression; in Spores of the Paleozoic of Byelorussia (Pripyat Depression), Byelorussian Scientific Research Institute of Geological Exploration, Collection of Scientific Works, p. 95-107, 140-146 (in Russian).

Ball, H.W. and Dineley, D.L.
1961: The Old Red Sandstone of Brown Clee Hill and the adjacent area. I. Stratigraphy; Bulletin of the British Museum (Natural History), Geology Series, v. 5, no. 7, p. 177-242.

Balme, B.E.
1962: Upper Devonian (Frasnian) spores from the Carnarvon Basin, Western Australia; The Palaeobotanist, v. 9, no. 1, 2, p. 1-10.

Balme, B.E. and Hennelly, J.P.F.
1956: Trilete sporomorphs from Australian Permian sediments; Australian Journal of Botany, v. 4, no. 3, p. 240-260.

Banks, H.P.
1980: Floral assemblages in the Siluro-Devonian; in Biostratigraphy of fossil plants, successional and paleoecological analyses, D.L. Dilcher and T.N. Taylor (eds.); Dowden, Hutchinson and Ross, Inc., Stroudsburg, Pennsylvania, p. 1-24.

Bassett, M.G., Cocks, L.R.M., Holland, C.H., Rickards, R.B., and Warren, P.T.
1975: The type Wenlock Series; Reports, Institute of Geological Sciences, no. 75/13, p. 1-19.

Bassett, M.G., Lawson, J.D., and White, D.E.
1982: The Downton Series as the fourth series of the Silurian System; Lethaia, v. 15, no. 1, p. 1-24.

Becker, G., Bless, M.J.M., Streel, M., and Thorez, J.
1974: Palynology and ostracode distribution in the Upper Devonian and basal Dinantian of Belgium and their dependence on sedimentary facies; Mededelingen Rijks Geologische Dienst, Nieuwe Serie, v. 25, no. 2, p. 9-99.

Beju, D.
1967: Quelques spores, acritarches et chitinozoaires d'âge Dévonien inférieur de la plate-forme Moesienne (Roumanie); Review of Palaeobotany and Palynology, v. 5, no. 1-4, p. 39-49.

Bless, M., Bouckaert, J., Conil, R., Dreesen, R., Groessens, E., Streel, M., and Thorez, J.
1974: Guidebook, Excursion D, International Symposium on Belgian Micropaleontological Limits; J. Bouckaert, and M. Streel (eds.); Brussels, p. 1-40.

Blieck, A.
1982: Les Hétérostracés (vertébrés, Agnathes) du Dévonien inférieur du nord de la France et du sud de la Belgique (Artois-Ardenne); Annales de la Société Géologique de Belgique, v. 105, p. 9-23.

Blieck, A. and Jahnke, H.
1980: Pteraspiden (Vertebrata, Heterostraci) aus den Unteren Siegener Schichten und ihre stratigraphischen Konsequenzen; Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, v. 159, no. 3, p. 360-378.

Bouckaert, J., Streel, M., and Thorez, J.
1970: Le Famennien et les couches de transition Dévonien-Carbonifère dans la vallée de l'Ourthe (sud de Liège, Synclinorium de Dinant); Colloque sur la Stratigraphie du Carbonifère, Congrès et Colloques Université de Liège, v. 55, p. 25-46.

Bouckaert, J., Streel, M., Thorez, J., and Mound, M.C.
1969: Biostratigraphic chart of the Famennian Stage (Upper Devonian) in the type localities of Belgium: A preliminary report; Journal of Paleontology, v. 43, no. 3, p. 727-734.

Brice, D.
1980: Givétien; in Les étages français et leurs stratotypes, C. Cavelier and J. Roger (eds.); Mémoire du Bureau de Recherches Géologiques et Minières, v. 109, p. 9-25.

Brice, D., Bultynck, P., Deunff, J., Loboziak, S. and Streel, M.
1979: Données biostratigraphiques nouvelles sur le Givétien et le Frasnien de Ferques (Boulonnais, France); Annales de la Société Géologique du Nord, v. 98, p. 325-344.

Brideaux, W.W., Chamney, T.P., Dunay, R.E., Fritz, W.H., Hopkins, W.S., Jeletzky, J.A., McGregor, D.C., Norford, B.S., Norris, A.W., Pedder, A.E.H., Sherrington, P.F., Sliter, W.V., Sweet, A.R., Uyeno, T.T., and Waterhouse, J.B.
1975: Biostratigraphic determinations of fossils from the subsurface of the districts of Franklin and Mackenzie; Geological Survey of Canada, Paper 74-39, p. 1-19.

Bultynck, P.
1970: Révision stratigraphique et paléontologique de la coupe type du Couvinien; Mémoires de l'Institut Géologique de l'Université de Louvain, v. 26, p. 1-152.

1982: Pridolian - lower Gedinnian conodont faunas from Naux (Neufchateau Synclinorium) and from Lievin (Dinant Synclinorium); Courier Forschungsinstitut Senckenberg, v. 55, p. 135-138.

Byvsheva, T.V.
1976: Zonal assemblage of spores from Devonian-Carboniferous boundary deposits of eastern regions of the Russian Plateau; in Results of Palynological Research on Precambrian, Paleozoic and Mesozoic of the the USSR, T.V. Byvsheva (ed.); Trudy Vsesoiuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Neftianogo Instituta, Kama Branch, no. 192, p. 67-93 (in Russian).

1980: Zonal spore assemblages of the Tournaisian Stage in the eastern regions of the Russian Platform; in Palynology in the USSR (1976-1980), E.D. Zaklinskaya, M.I. Neishtadt, A.M. Medvedeva, and L.V. Rovnina (eds.); "Nauka", Moscow, p. 53-55 (in Russian).

Chaloner, W.G.
1959: Devonian megaspores from Arctic Canada; Palaeontology, v. 1, pt. 4, p. 321-332.

Chi, B.I. and Hills, L.V.
1974: Stratigraphic and paleoenvironmental significance of Upper Devonian megaspores, type section of the Imperial Formation, Northwest Territories, Canada; in Canadian Arctic Geology; Proceedings of Symposium on Geology of Canadian Arctic, Saskatoon, J.D. Aitken and D.J. Glass (eds.), p. 241-257.

1976: Biostratigraphy and taxonomy of Devonian megaspores, Arctic Canada; Bulletin of Canadian Petroleum Geology, v. 24, no. 4, p. 640-818.

Chibrikova, E.V.
1959: Spores from the Devonian and older rocks of Bashkiria; Academy of Sciences of USSR, Bashkirian Branch, Data on Paleontology and Stratigraphy of Devonian and Older Deposits of Bashkiria, p. 3-116 (in Russian).

1962: Spores of Devonian terrigenous deposits of western Bashkiria and the western slopes of the southern Urals; in Brachiopods, ostracods and spores of the Middle and Upper Devonian of Bashkiria; Academy of Sciences of the USSR, Bashkirian Branch, Institute of Mining Geology; p. 353-476 (in Russian).

1972: Plant microfossils from the South Urals and Cisuralia; Academy of Sciences of the USSR, Bashkirian Branch, Institute of Geology, "Nauka", Moscow, p. 1-222 (in Russian).

1977: Stratigraphy of Devonian and older Paleozoic deposits of the southern Urals and Priuralia (on plant microfossils); Academy of Sciences of the USSR, Bashkirian Branch, Institute of Geology, "Nauka", Moscow, p. 1-191 (in Russian).

1978: The Lower-Middle Devonian boundary on the western slope of the southern Urals based on the evidence of spore studies; Problems of stratigraphy of the Paleozoic (Devonian, Carboniferous), Academy of Sciences of the USSR, Interdepartmental Stratigraphic Committee of the USSR, Transactions, v. 6, p. 107-112 (in Russian).

Chibrikova, E.V. (cont.)
1983: Palynological substantiation of the boundaries and differentiation of the Middle Devonian; in The Middle Devonian of the USSR, its boundaries and differentiation into stages, M.A. Kamaletdinov (ed.); Academy of Sciences of the USSR, Bashkirian Branch, Institute of Geology, Ufa, p. 25-26 (in Russian).

Clayton, G., Coquel, R., Doubinger, J., Gueinn, K.J., Loboziak, S., Owens, B., and Streel, M.
1977: Carboniferous miospores of western Europe: illustration and zonation; Mededelingen Rijks Geologische Dienst, v. 29, 71 p.

Clayton, G. and Graham, J.R.
1974: Miospore assemblages from the Devonian Sherkin Formation of southwest County Cork, Republic of Ireland; Pollen et Spores, v. 16, no. 4, p. 565-588.

Clayton, G., Higgs, K., Gueinn, K.J., and van Gelder, A.
1974: Palynological correlations in the Cork Beds (Upper Devonian - ?Upper Carboniferous) of Southern Ireland; Proceedings of Royal Irish Academy, v. 74, sec. B, no. 10, p. 145-155.

Clayton, G., Higgs, K., Keegan, J.B., and Sevastopulo, G.D.
1978: Correlation of the palynological zonation of the Dinantian of the British Isles; Palinología, número extraordinario 1, p. 137-148.

Clendening, J.A., Eames, L.E., and Wood, G.D.
1980: *Retusotriletes phillipsii* n. sp., a potential Upper Devonian guide palynomorph; Palynology, v. 4, p. 15-22.

Cocks, L.R.M., Holland, C.H., Rickards, R.B., and Strachan, I.
1971: A correlation of Silurian rocks in the British Isles; Geological Society of London, Special Report No. 1, Journal of the Geological Society, v. 127, p. 103-106.

Cocks, L.R.M., Lane, P.D., Rickards, R.B., Temple, J.T., and Woodcock, N.H.
1983: The Llandovery area as the type for the first Series of the Silurian System; International Union of Geological Sciences, Subcommission on Silurian Stratigraphy, p. 1-30.

Combaz, A.
1967: Un microbios du trémadocien dans un sondage d'Hassi-Messaoud; Actes de la Société Linnéenne de Bordeaux, v. 104, sér. B, no. 29, p. 1-26.

Conil, R. and Lys, M.
1980: Strunien; in Les étages français et leurs stratotypes, C. Cavelier and J. Roger (eds.); Mémoire du Bureau de Recherches Géologiques et Minières, no. 109, p. 26-35.

Cooper, B.J.
1980: Toward an improved Silurian conodont biostratigraphy; Lethaia, v. 13, p. 209-227.

Cramer, F.H.
1966a: Hoegispheres and other microfossils *incertae sedis* of the San Pedro Formation (Siluro-Devonian boundary) near Valporquero, León, NW Spain; Notas y Comunicaciones del Instituto Geológico y Minero de España, no. 86, p. 75-94.

Cramer, F.H. (cont.)
1966b: Palynomorphs from the Siluro-Devonian boundary in NW Spain; Notas y Comunicaciónes del Instituto Geológico y Minero de España, no. 85, p. 71-82.

1967: Palynology of Silurian and Devonian rocks in northwest Spain; Boletín del Instituto Geológico y Minero de España, v. 77, p. 223-286.

D'Erceville, M.A.
1979: Les spores des formations Siluro-Dévoniennes de la coupe de Saint-Pierre-sur-Erve (Synclinorium médian armoricain); Palaeontographica, Abteilung B, v. 171, p. 79-121.

Dineley, D.L. and Williams, B.P.J.
1968: The Devonian continental rocks of the lower Restigouche River, Quebec; Canadian Journal of Earth Sciences, v. 5, p. 945-953.

Dolby, G. and Neves, R.
1970: Palynological evidence concerning the Devonian-Carboniferous boundary in the Mendips, England; Compte Rendu, Sixième Congrès International de Stratigraphie et de Géologie du Carbonifère, Ernest van Aelst Publishing Company, Netherlands, v. 2, p. 631-646.

Edwards, D.
1968: A new plant from the Lower Old Red Sandstone of south Wales; Palaeontology, v. 11, pt. 5, p. 683-690.

Egorova, V.L.
1974: New species of spores from the Middle Devonian of Kiubishev District; Geology and Geochemistry, Kiubishev, 1974, p. 31-35 (in Russian).

Embry, A.F. and Klovan, J.E.
1976: The Middle-Upper Devonian clastic wedge of the Franklinian geosyncline; Bulletin of Canadian Petroleum Geology, v. 24, no. 4, p. 485-639.

Errera, M., Mamet, B., and Sartenaer, P.
1972: Le Calcaire de Givet et le Givetien à Givet; Institut Royal des Sciences Naturelles de Belgique, Bulletin, Sciences de la Terre, v. 48, no. 1, p. 1-59.

Evans, P.R.
1970: Revision of the miospore genera *Perotrilites* Erdtm. ex Couper 1953 and *Diaphanospora* Balme and Hassell 1962; Palaeontological Papers, 1968, Australia, Bureau of Mineral Resources, Geology and Geophysics, Bulletin no. 116, p. 65-74.

Fomina, L.S.
1968: Spore assemblages of the terrigenous Devonian in northern areas of the Volga-Urals region; in Paleopalynological method in stratigraphy, All-Union Scientific Research Institute of Geology (VSEGEI), Leningrad, p. 5-10 (in Russian).

Ford, J.H.
1972: Palynology of some Upper Silurian and Lower Old Red Sandstone sediments in the Midland Valley of Scotland; unpublished Ph.D. thesis, King's College, London.

Fuchs, G.
1974: Das Unterdevon am Ostrand der Eifeler Nordsüd-Zone; Beiträge zur naturkundlichen Forschung in Südwestdeutschland, Beiheft 2, p. 3-163.

Gao Lianda
1981: Devonian spore assemblages of China; Review of Palaeobotany and Palynology, v. 34, p. 11-23.

1983: Discovery of Late Devonian spore assemblage from Nyalam County, Xizang (Tibet) and its stratigraphic significance; 8 CGOXP Editorial Committee, Ministry of Geology and Mineral Resources, People's Republic of China, Geological Publishing House, Beijing, p. 183-218 (in Chinese).

Gao Lianda and Hou Jingpeng
1975: Early and Middle Devonian spore assemblages from Dushan and Guyun of Kueichow and their stratigraphical significance; Professional Papers of Stratigraphy and Palaeontology, no. 1, Geological Press, Beijing, p. 170-232 (in Chinese).

Gensel, P.G.
1979: Two *Psilophyton* species from the Lower Devonian of eastern Canada with a discussion of morphological variation within the genus; Palaeontographica, Abteilung B, v. 168, p. 81-99.

Godefroid, J., Stainer, P., and Steemans, P.
1982: Gedinnian and Siegenian lithostratigraphy and biostratigraphy of Belgium; in Field meeting on Lower and Middle Devonian stages in the Ardenno-Rhenish type area, Guidebook, G. Plodowski, R. Werner, and W. Ziegler (eds.); Subcommission on Devonian Stratigraphy, Frankfurt am Main, p. 3-79.

Golubtsov, V.K. (editor)
1978: Stratigraphical and paleontological research in Byelorussia; Byelorussian Scientific Research Geological Exploration Institute, "Nauk i Tekhnika", Minsk, p. 1-248 (in Russian).

Gray, J. and Boucot, A.J.
1971: Early Silurian spore tetrads from New York: earliest new world evidence for vascular plants?; Science, v. 173, p. 918-921.

Gray, J., Massa, D., and Boucot, A.J.
1982: Caradocian land plant microfossils from Libya; Geology, v. 10, no. 4, p. 197-201.

Grishina, T.S. and Klenina, L.N.
1981: Correlation of marine and continental middle-upper Paleozoic deposits of eastern Kazakhstan; in Life on ancient continents, its growth and development, I.V. Vasiliev and L.I. Khozatskii (eds.); Transactions, 23rd session of the All-Union Paleontological Society, Leningrad, "Nauka", p. 51-57 (in Russian).

Guennel, G.K.
1963: Devonian spores in a Middle Silurian reef; Grana Palynologica, v. 4, no. 2, p. 243-261.

Hacquebard, P.A.
1957: Plant spores from the Horton Group (Mississippian) of Nova Scotia; Micropaleontology, v. 3, no. 4, p. 301-324.

Hancock, J.M.
1977: The historic development of concepts of biostratigraphical correlation; in Concepts and methods of biostratigraphy, E.G. Kauffman and J.E. Hazel (eds.); Dowden, Hutchinson and Ross, Stroudsburg, Pennsylvania, p. 3-22.

Hassan, A.
1982: Palynology, stratigraphy and provenance of the Lower Old Red Sandstone of the Brecon Beacons (Powys) and Black Mountains (Gwent and Powys), South Wales; unpublished Ph.D. thesis, University of London, King's College, v. 1, 440 p.; v. 2, text-figs. and plates.

Hedberg, H.D. (ed.)
1976: International stratigraphic guide; John Wiley and Sons, New York, 200 p.

Higgs, K.
1975: Upper Devonian and Lower Carboniferous miospore assemblages from Hook Head, County Wexford, Ireland; Micropaleontology, v. 21, no. 4, p. 393-419.

Higgs, K. and Russell, K.J.
1981: Upper Devonian microfloras from southeast Iveragh, County Kerry, Ireland; Geological Survey of Ireland, Bulletin, v. 3, no. 1, p. 17-50.

Hill, P.J.
1974: Stratigraphic palynology of acritarchs from the type areas of the Llandovery and the Welsh Borderland; Review of Palaeobotany and Palynology, v. 18, p. 11-23.

Hoffmeister, W.S.
1959: Lower Silurian plant spores from Libya; Micropaleontology, v. 5, no. 3, p. 331-334.

Holland, C.H.
1981: Devonian; in A geology of Ireland, C.H. Holland (ed.); Scottish Academic Press, Edinburgh, p. 121-146.

Holland, C.H., Audley-Charles, M.G., Bassett, M.G., Cowie, J.W. Curry, D., Fitch, F.J., Hancock, J.M., House, M.R., Ingham, J.K., Kent, P.E., Morton, N., Ramsbottom, W.H.C., Rawson, P.F., Smith, D.B., Stubblefield, C.J., Torrens, H.S., Wallace, P., and Woodland, A.W.
1978: A guide to stratigraphical procedure; Geological Society of London, Special Report no. 11, p. 1-18.

Holland, C.H., Lawson, J.D. and Walmsley, V.G.
1963: The Silurian rocks of the Ludlow District, Shropshire; Bulletin of the British Museum (Natural History), Geology Series, v. 8, no. 3, p. 93-171.

Holland, C.H. and Richardson, J.B.
1977: The British Isles; in The Silurian-Devonian boundary, A. Martinsson, (ed.); International Union of Geological Sciences, Series A, Stuttgart, no. 5, p. 35-44.

Hou Jingpeng
1982: Some spore assemblages of the Devonian-Carboniferous transitional beds in the Xikuangshan region, central Hunan; Bulletin of the Institute of Geology, Chinese Academy of Geological Sciences, Beijing, v. 1982, no. 5, p. 81-95 (in Chinese).

House, M.R., Richardson, J.B., Chaloner, W.G., Allen, J.R.L., Holland, C.H., and Westoll, T.S.
1977: A correlation of Devonian rocks in the British Isles; Geological Society of London, Special Report no. 8, p. 1-110.

House, M.R. and Ziegler, W.
1977: The goniatite and conodont sequences in the early Upper Devonian at Adorf, Germany; Geologica et Paleontologica, v. 11, p. 69-108.

Hughes, N.F.
1975: The challenge of abundance in palynomorphs; Geoscience and Man, v. 11, p. 141-144.

Jaeger, H.
1979: Devonian Graptolithina; in The Devonian System, M.R. House, C.T. Scrutton, and M.G. Bassett (eds.); Palaeontological Association, London, Special Papers in Paleontology, no. 23, p. 335-339.

Jardiné, S. and Yapaudjian, L.
1968: Lithostratigraphie et palynologie du Dévonien-Gothlandien gréseux du Bassin de Polignac (Sahara); Revue de l'Institut Français du Pétrole, v. 23, no. 4, p. 439-469.

Johnson, J.G.
1979: Intent and reality in biostratigraphic zonation; Journal of Paleontology, v. 53, no. 4, p. 931-942.

Johnson, N.G.
1984: Early Silurian palynomorphs from the Tuscarora Formation in central Pennsylvania and their paleobotanical and geological significance; unpublished M.Sc. thesis, Pennsylvania State University, p. 1-97.

Kaiser, H.
1971: Die Oberdevon-Flora der Bäreninsel. 4. Mikroflora der Misery-Serie und der Flözleeren Sandstein-Serie; Palaeontographica, Abteilung B, v. 135, p. 127-164.

Kaljo, D.
1978: The Downtonian or Pridolian from the point of view of the Baltic Silurian; Eesti NSV Teaduste Akadeemia Toimetised, Geoloogia, v. 27, no. 1, p. 5-10.

Karathanasopoulos, S.
1975: Die Sporenvergesellschaftungen in den Dachschiefern des Hunsrücks (Rheinisches Schiefergebirge, Deutschland) und ihre Aussage zur Stratigraphie; unpublished doctoral dissertation, University of Mainz, 96 p.

Kedo, G.I.
1955: Spores of the Middle Devonian of the northeastern Byelorussian SSR; Academy of Sciences of BSSR, Institute of Geological Sciences, Paleontology and Stratigraphy of BSSR, Sbornik 1, p. 5-59 (in Russian).

1963: Spores of the Tournaisian stage of the Pripyat Depression and their stratigraphic significance; Academy of Sciences of BSSR, Institute of Geological Sciences, Paleontology and Stratigraphy of BSSR, Sbornik 4, p. 3-121 (in Russian).

Kedo, G.I., Nazarenko, A.M., Nekryata, N.S., Raskatova, I.G., Sennova, V.F., and Chibrikova, E.V.
1971: New species of spores from Famennian deposits of the Pripyat Depression, central regions of the Russian Platform, Volga-Ural oil and gas province and Timan; BelNIGRI, Palynological Research in Byelorussia and other regions of USSR, p. 172-205 (in Russian).

Kedo, G.I. and Obukhovskaya, T.G.
1981: Spores, Middle Devonian of the Baltic region and northeastern White Russia; in Devonian and Carboniferous of the Baltic regions, V.S. Sorokin, L.A. Liarskaya and L.S. Savvaitova, (eds.); Riga, "Zinatne", p. 419-436 (in Russian).

Kemp, E.M.
1972: Lower Devonian palynomorphs from the Horlick Formation, Ohio Range, Antarctica; Palaeontographica, Abteilung B, v. 139 p. 105-124.

Keppie, J.D., Giles, P.S., and Boehner, R.C.
1978: Some Middle Devonian to Lower Carboniferous rocks of Cape George, Nova Scotia; Nova Scotia Department of Mines, Paper 78-4, p. 1-37.

Kerr, J.W., McGregor, D.C., and McLaren, D.J.
1965: An unconformity between Middle and Upper Devonian rocks of Bathurst Island, with comments on Upper Devonian faunas and microfloras of the Parry Islands; Bulletin of Canadian Petroleum Geology, v. 13, no. 3, p. 409-431.

Kirchgasser, W.T. and House, M.R.
1981: Upper Devonian goniatite biostratigraphy; in Devonian biostratigraphy of New York, Part I, text, W.A. Oliver, Jr. and G. Klapper (eds.); Subcommission on Devonian Stratigraphy, Washington, D.C., p. 39-55.

Klapper, G. and Johnson, J.G.
1980: Endemism and dispersal of Devonian conodonts; Journal of Paleontology, v. 54, no. 2, p. 400-455.

Kushnareva, T.I., Khalymbadzha, V.G. and Busygina, Yu.N.
1978: Biostratigraphic zonation of the Domanik Formation in the stratotype section; Soviet Geology, v. 1978, no. 1, p. 60-71 (in Russian).

Lanninger, E.P.
1968: Sporen-Gesellschaften aus dem Ems der SW-Eifel; Palaeontographica, Abteilung B, v. 122, p. 95-170.

Laufeld, S.
1974: Reference localities for palaeontology and geology in the Silurian of Gotland; Sveriges Geologiska Undersokning, Series C, no. 705, p. 1-172.

Lawson, J.D.
1954: The Silurian succession at Gorsley (Herefordshire); Geological Magazine, v. 91, no. 3, p. 227-237.

Loboziak, S. and Streel, M.
1980: Miospores in Givetian to lower Frasnian sediments dated by conodonts from the Boulonnais, France; Review of Palaeobotany and Palynology, v. 29, p. 285-299.

Loboziak, S. and Streel, M. (cont.)
1981: Miospores in middle-upper Frasnian to Famennian sediments partly dated by conodonts (Boulonnais, France); Review of Palaeobotany and Palynology, v. 34, p. 49-66.

Loboziak, S., Streel, M. and Vanguestaine, M.
1983: Miospores et acritarches de la Formation d'Hydrequent (Frasnien supérieur à Famennien inférieur), Boulonnais, France; Annales de la Société Géologique de Belgique, v. 106, p. 173-183.

Lu Lichang
1980a: Devonian miospores from the Longhuashan section in Zhanyi of Yunnan and their stratigraphic significance; Memoirs of Nanjing Institute of Geology and Palaeontology, Academia Sinica, no. 14, p. 1-62 (in Chinese).

1980b: On the occurrence of *Archaeoperisaccus* in East Yunnan; Acta Palaeontologica Sinica, v. 19, no. 6, p. 500-502 (in Chinese).

Mamet, B.L.
1977: Foraminiferal zonation of the Lower Carboniferous: methods and stratigraphic implications; in Concepts and methods of biostratigraphy, E.G. Kauffman and J.E. Hazel (eds.); Dowden, Hutchinson and Ross, Stroudsburg, Pennsylvania, p. 445-462.

Marshall, J.E.A. and Allen, K.C.
1982: Devonian miospore assemblages from Fair Isle, Shetland; Palaeontology, v. 25, pt. 2, p. 277-312.

Martin, F.
1978: Lower Paleozoic Chitinozoa and Acritarcha from Newfoundland; in Current Research, Part B, Geological Survey of Canada, Paper 78-1B, p. 73-81.

Martinsson, A.
1977: Baltoscandia; in The Silurian-Devonian boundary, A. Martinsson (ed.); International Union of Geological Sciences, Series A, Stuttgart, no. 5, p. 45-51.

Maslov, V.A., Abramova, A.N., Baryshev, V.N., Chibrikova, E.V., Zhavoronkova, R.A., and Garifullina, A.A.
1983: Stratigraphy of the Lower and Middle Devonian of the western slope of the South Urals; in The Middle Devonian of the USSR, its boundaries and differentiation into stages, M.A. Kamaletdinov (ed.); Academy of Sciences of the USSR, Bashkirian Branch, Institute of Geology, Ufa, p. 7-9 (in Russian).

Massa, D.
1980: A stratigraphic contribution to the Palaeozoic of the southern basins of Libya. Part I. General framework of the Murzuq and Ghadamis basins. Part II. The Murzuq Basin; Bellini, E. and Massa, D., in The geology of Libya, M.J. Salem and M.I. Busrewil (eds.); Academic Press, London, v. 1, p. 6-32.

Massa, D. and Moreau-Benoit, A.
1976: Essai de synthèse stratigraphique et palynologique du système Dévonien en Libye Occidentale; Revue de l'Institut Français du Pétrole, v. 31, no. 2, p. 287-333.

McCracken, A.D. and Barnes, C.R.
1981: Conodont biostratigraphy and paleoecology of the Ellis Bay Formation, Anticosti Island, Quebec, with special reference to Late Ordovician-Early Silurian chronostratigraphy and the systemic boundary; Geological Survey of Canada, Bulletin 329, p. 51-134.

McGregor, D.C.
1960: Devonian spores from Melville Island, Canadian Arctic Archipelago; Palaeontology, v. 3, pt. 1, p. 26-44.

1961: Spores with proximal radial pattern from the Devonian of Canada; Geological Survey of Canada, Bulletin 76, 11 p.

1964: Devonian miospores from the Ghost River Formation, Alberta; Geological Survey of Canada, Bulletin 109, 31 p.

1967: Composition and range of some Devonian spore assemblages of Canada; Review of Palaeobotany and Palynology, v. 1, p. 173-183.

1969: Devonian plant fossils of the genera *Kryshtofovichia, Nikitinsporites,* and *Archaeoperisaccus;* in Contributions to Canadian Paleontology, Norford, B.S. et al., Geological Survey of Canada, Bulletin 182, p. 91-106.

1970: *Hymenozonotriletes lepidophytus* Kedo and associated spores from the Devonian of Canada; Colloque sur la Stratigraphie du Carbonifère, Les Congrès et Colloques de l'Université de Liège, v. 55, p. 315-326.

1973: Lower and Middle Devonian spores of eastern Gaspé, Canada. I. Systematics; Palaeontographica, Abteilung B, v. 142, p. 1-77.

1974: Early Devonian spores from central Ellesmere Island, Canadian Arctic; Canadian Journal of Earth Sciences, v. 11, p. 70-78.

1977: Lower and Middle Devonian spores of eastern Gaspé, Canada. II. Biostratigraphic significance; Palaeontographica, Abteilung B, v. 163, p. 111-142.

1979a: Devonian spores from the Barrandian region of Czechoslovakia and their significance for interfacies correlation; in Current Research, Part B, Geological Survey of Canada, Paper 79-1B, p. 189-197.

1979b: Spores in Devonian stratigraphical correlation; in The Devonian System, M.R. House, C.T. Scrutton and M.G. Bassett (eds.); Palaeontological Association, Special Paper 23, p. 163-184.

1979c: Devonian miospores of North America; Palynology, v. 3, p. 31-52.

1981: Spores and the Middle-Upper Devonian boundary; Review of Palaeobotany and Palynology, v. 34, p. 25-47.

1984: Late Silurian and Devonian spores from Bolivia; Academia Nacional de Ciencias, Córdoba, Miscelanea no. 69, 57 p.

McGregor, D.C. and Camfield, M.
1976: Upper Silurian(?) to Middle Devonian spores of the Moose River Basin, Ontario; Geological Survey of Canada, Bulletin 263, 63 p.

1982: Middle Devonian miospores from the Cape De Bray, Weatherall, and Hecla Bay formations of northeastern Melville Island, Canadian Arctic; Geological Survey of Canada, Bulletin 348, 105 p.

McGregor, D.C. and Narbonne, G.M.
1978: Upper Silurian trilete spores and other microfossils from the Read Bay Formation, Cornwallis Island, Canadian Arctic; Canadian Journal of Earth Sciences, v. 15, p. 1292-1303.

McGregor, D.C. and Owens, B.
1966: Illustrations of Canadian fossils: Devonian spores of eastern and northern Canada; Geological Survey of Canada, Paper 66-30, 66 p.

McGregor, D.C. and Uyeno, T.T.
1972: Devonian spores and conodonts of Melville and Bathurst islands, District of Franklin; Geological Survey of Canada, Paper 71-13, 37 p.

McKerrow, W.S., Lambert, R. St. J., and Cocks, L.R.M.
1985: The Ordovician, Silurian and Devonian periods; in The geological time scale, N.J. Snelling (ed.); Geological Society of London, Special Publication no. 15, p. 73-80.

Medianik, S.I.
1980: Spores of the genera *Archaeoperisaccus* Naumova and *Archaeomonoletes* gen. n. and their stratigraphic meaning in the Timan-Pechora provinces; Vestnik Moscow University, Geology, v. 1980, no. 5, p. 93-96 (in Russian).

1982: Sporangia of lower Frasnian *Archaeopteris* from southern Timan; Paleontological Journal, v. 1982, no. 2, p. 121-127 (in Russian).

Miall, A.D.
1976: Proterozoic and Paleozoic geology of Banks Island, Arctic Canada; Geological Survey of Canada, Bulletin 258, 77 p.

Mikhailova, T.E. and Fradkina, A.F.
1966: Palynologic characteristics of Devonian-Lower Carboniferous deposits of the Kempendyai structural-prospecting borehole section; Academy of Sciences of USSR, Siberian Branch, Institute of Geology and Geophysics, Palynology of Siberia, 2nd International Conference on Palynology, p. 70-74 (in Russian).

Miller, M.E. and Eames, L.E.
1982: Palynomorphs from the Silurian Medina Group (lower Llandovery) of the Niagara Gorge, Lewiston, New York, U.S.A.; Palynology, v. 6, p. 221-254.

Moreau-Benoit, A.
1976: Les spores et débris végétaux; in Les Schistes et Calcaires Eodévoniens de Saint-Cénére (Massif Armoricain France); Mémoires de la Société géologique et minéralogique de Bretagne, no. 19, p. 27-57.

Moreau-Benoit, A. (cont.)
1979: Les spores du Dévonien de Libye (première partie); Cahiers de Micropaléontologie, Centre National de la Recherche Scientifique, Paris, v. 1979, no. 4, 58 p.

1980: Les spores du Dévonien de Libye (deuxième partie); Cahiers de Micropaléontologie, Centre National de la Recherche Scientifique, Paris, v. 1980, no. 1, 53 p.

Mortimer, M.G. and Chaloner, W.G.
1967: Devonian megaspores from the Wyboston borehole, Bedfordshire, England; Palaeontology, v. 10, pt. 2, p. 189-213.

1972: The palynology of concealed Devonian rocks of southern England; Geological Survey of Great Britain, Bulletin 39, p. 1-56.

Murphy, M.A. and Berry, W.B.N.
1983: Early Devonian conodont-graptolite collation and correlations with brachiopod and coral zones, central Nevada; American Association of Petroleum Geologists, Bulletin, v. 67, no. 3, p. 371-379.

Naumova, S.N.
1953: Spore-pollen assemblages of the Upper Devonian of the Russian Platform and their stratigraphic significance; Transactions of Institute of Geological Sciences, Academy of Sciences of the USSR, no. 143 (Geological Series no. 60), 204 p. (in Russian).

Naylor, D.
1975: Upper Devonian-Lower Carboniferous stratigraphy along the south coast of Dunmanus Bay, Co. Cork; Royal Irish Academy, Proceedings, Section B, v. 75, p. 317-337.

Nazarenko, A.M.
1965: A characterization of the spore assemblages of the Middle Devonian beds of the province of Volgograd; Proceedings of the Volgograd Scientific-Research Institute of the Petroleum and Gas Industry (VNIING), no. 3, "Nedra", Leningrad, p. 39-47 (in Russian).

1974: Famennian assemblages of plant microfossils from the Volgograd region of the Volga and their changes related to the lithological composition of the enclosing rocks; Palynology of Proterophyte and Paleophyte, Proceedings of III International Palynological Conference, S.N. Naumova (ed.); Moscow, "Nauka", p. 82-85 (in Russian).

Nazarenko, A.M., Kedo, G.I., Araslanova, R.M., Zhdanova, L.R., Lomaeva, E.T., Raskatova, L.G., Sennova, V.F., Sergeeva, L.A., Umnova, V.T., and Shishova, G.M.
1971: New species of spores from the Frasnian Stage of the Russian Platform and Altai; BelNIGRI, Palynological Research in Byelorussia and other Regions of USSR, Minsk, p. 155-172 (in Russian).

Norris, A.W.
1968: Reconnaissance Devonian stratigraphy of northern Yukon Territory and northwestern District of Mackenzie; Geological Survey of Canada, Paper 67-53, 287 p.

North American Commission on Stratigraphic Nomenclature
1983: North American stratigraphic code; American Association of Petroleum Geologists, Bulletin, v. 67, no. 5, p. 841-875.

Oliver, W.A., Jr. and Klapper, G. (eds.)
1981: Devonian biostratigraphy of New York, Part I, Text; International Union of Geological Sciences, Subcommission on Devonian Stratigraphy, Washington, D.C., p. 1-106.

Owens, B.
1971: Miospores from the Middle and early Upper Devonian rocks of the western Queen Elizabeth Islands, Arctic Archipelago; Geological Survey of Canada, Paper 70-38, 157 p.

Paproth, E., Conil, R., Bless, M.J.M., Boonen, P., Carpentier, N., Coen, M., Delcambre, B., Deprijck, Ch., Deuzon, S., Dreesen, R., Groessens, E., Hance, L., Hennebert, M., Hibo, D., Hahn, G. & R., Hislaire, O., Kasig, W., Laloux, M., Lauwers, A., Lees, A., Lys, M., Op de Beek, K., Overlau, P., Pirlet, H., Poty, E., Ramsbottom, W., Streel, M., Swennen, R., Thorez, J., Vanguestaine, M., Van Steenwinkel, M., and Vieslet, J.L.
1983: Bio- and lithostratigraphic subdivisions of the Dinantian in Belgium, a review; Annales de la Société Géologique de Belgique, v. 106, p. 185-239.

Paproth, E. and Streel, M.
1970: Corrélations biostratigraphiques près de la limite Dévonien/Carbonifère entre les faciès littoraux ardennais et les faciès bathyaux rhénans; Colloque sur la Stratigraphie Carbonifère, Congrès et Colloques de l'Université de Liège, v. 55, p. 365-398.

Pashkevich, N.G.
1964: New Devonian species of *Archaeoperisaccus* (Gymnospermae) from Northern Timan; Palaeontological Journal, v. 1964, no. 4, p. 126-129 (in Russian).

1971: Comparative character of Famennian spore assemblages of the region of the Kempendyai dislocation (Siberian Platform) and Northern Timan (Russian Platform); Palynological Characteristics of Paleozoic, Mesozoic and Cenozoic Deposits of Yakutia; Yakutsk, p. 40-45 (in Russian).

Paulus, B.
1961: Das Urfttal-Profil in der Sötenicher Eifelkalkmulde (Devon); Der Aufschluss, H. "Eifel", Heidelberg, p. 26-40.

Pepper, J.F. and de Witt, W.
1950: Stratigraphy of the Upper Devonian Wiscoy Sandstone and the equivalent Hanover Shale in western and central New York; United States Geological Survey, Oil and Gas Preliminary Chart 37.

Playford, G.
1963: Lower Carboniferous microfloras of Spitsbergen-Part Two; Palaeontology, v. 5, part 4, p. 619-678.

1964: Miospores from the Mississippian Horton Group of eastern Canada; Geological Survey of Canada, Bulletin 107, 47 p.

1971: Lower Carboniferous spores from the Bonaparte Gulf Basin, Western Australia and Northern Territory; Australia, Bureau of Mineral Resources, Geology and Geophysics, Bulletin 115, p. 1-102.

1976: Plant microfossils from the Upper Devonian and Lower Carboniferous of the Canning Basin, Western Australia; Palaeontographica, Abteilung B, v. 158, p. 1-71.

1983: The Devonian miospore genus *Geminospora* Balme 1962: a reappraisal based upon topotypic *G. lemurata* (type species); Memoirs of the Association of Australasian Palaeontologists, v. 1, p. 311-325.

Pratt, L.M., Phillips, T.L., and Dennison, J.M.
1978: Evidence of non-vascular land plants from the Early Silurian (Llandoverian) of Virginia, U.S.A.; Review of Palaeobotany and Palynology, v. 25, p. 121-149.

Pychova, N.G.
1960: Spore-pollen assemblages of the terrigenous part of the productive Devonian in the Tatar ASSR and their significance for stratigraphy; Transactions of the All-Union Oil and Gas Scientific Research Institute (VNII), no. 23, The Geology of Oil Deposits, p. 37-48 (in Russian).

Raskatova, L.G.
1973: Palynological description of the Famennian deposits in the central regions of the Russian Platform; Voronezh University Press, Voronezh, 176 p. (in Russian).

1974: A detailed stratigraphic subdivision of the Devonian of the Central Devonian Field in the light of palynological evidence; Palynology of Proterophyte and Paleophyte, Proceedings III International Palynological Conference, S.N. Naumova (ed.); Moscow, "Nauka", p. 67-71 (in Russian).

Raskatova, L.G., Shishova, G.M., Kholmovaya, R.S., Neberikutina, L.N., and Raskatova, M.G.
1983: Methods of correlation of Frasnian deposits of the Central Devonian Field and the Timan-Pechora Province; in Stratigraphy and correlation of sediments using palynological methods, 4th All-Union Palynological Conference, Tiumen, G.N. Papulov (ed.); Academy of Sciences of USSR, Sverdlovsk, p. 50-52 (in Russian).

Rice, R.J.
1982: Sedimentology of the late Middle and Upper Devonian Okse Bay Group, southwestern Ellesmere Island: a progress report; in Current Research, Part A, Geological Survey of Canada, Paper 82-1A, p. 79-82.

Richards, B.C.
in press: Uppermost Devonian and Lower Carboniferous stratigraphy, sedimentation and diagenesis of southwestern District of Mackenzie and southeastern Yukon Territory (NTS 95B, C, F and G); Geological Survey of Canada, Bulletin.

Richardson, J.B.
1960: Spores from the Middle Old Red Sandstone of Cromarty, Scotland; Palaeontology, v. 3, pt. 1, p. 45-63.

1962: Spores with bifurcate processes from the Middle Old Red Sandstone of Scotland; Palaeontology, v. 5, pt. 2, p. 171-194.

1965: Middle Old Red Sandstone spore assemblages from the Orcadian Basin, northeast Scotland; Palaeontology, v. 7, pt. 4, p. 559-605.

1967: Some British Lower Devonian spore assemblages and their stratigraphic significance; Review of Palaeobotany and Palynology, v. 1, no. 1-4, p. 111-129.

1969: Devonian spores; in Aspects of Palynology, R.H. Tschudy and R.A. Scott (eds.); Wiley-Interscience, p. 193-222.

1974: The stratigraphic utilization of some Silurian and Devonian miospore species in the northern hemisphere: an attempt at a synthesis; International Symposium on Belgian Micropaleontological Limits, Namur 1974, Publication no. 9, p. 1-13.

Richardson, J.B., Ford, J.H., and Parker, F.
1984: Miospores, correlation and age of some Scottish Lower Old Red Sandstone sediments from the Strathmore region (Fife and Angus); Journal of Micropalaeontology, v. 3, no. 2, p. 109-124.

Richardson, J.B. and Ioannides, N.S.
1973: Silurian palynomorphs from the Tanezzuft and Acacus formations, Tripolitania, North Africa; Micropaleontology, v. 19, p. 257-307.

1979: *Emphanisporites splendens*, new name for *Emphanisporites pseudoerraticus* Richardson and Ioannides, 1973 (preoccupied); Micropaleontology, v. 25, p. 111.

Richardson, J.B. and Lister, T.R.
1969: Upper Silurian and Lower Devonian spore assemblages from the Welsh Borderland and South Wales; Palaeontology, v. 12, pt. 2, p. 201-252.

Richardson, J.B. and Rasul, S.M.
1978: Palynomorphs in Lower Devonian sediments from the Apley Barn Borehole, southern England; Pollen et Spores, v. 20, no. 3, p. 423-462.

Richardson, J.B., Rasul, S.M., and Al-Ameri, T.
1981: Acritarchs, miospores and correlation of the Ludlovian-Dowtonian and Silurian-Devonian boundaries; Review of Palaeobotany and Palynology, v. 34, p. 209-224.

Richardson, J.B., Streel, M., Hassan, A., and Steemans, Ph.
1982: A new spore assemblage to correlate between the Breconian (British Isles) and the Gedinnian (Belgium); Annales de la Société Géologique de Belgique, v. 105, p. 135-143.

Rickard, L.V.
1975: Correlation of the Silurian and Devonian rocks in New York State; New York State Museum and Science Service, Map and Chart Series No. 24, University of the State of New York and State Education Department, Albany, 16 p.

Riegel, W.
1968: Die Mitteldevon-Flora von Lindlar (Rheinland). 2. Sporae dispersae; Palaeontographica, Abteilung B, v. 123, p. 76-96.

1973: Sporenformen aus den Heisdorf-, Lauch- und Nohn-Schichten (Emsium und Eifelium) der Eifel, Rheinland; Palaeontographica, Abteilung B, v. 142, p. 78-104.

1975: Die dispersen Sporen der Ems-, Eifel- und Givet-Stufe der Eifel (Rheinisches Schiefergebirge) und ihre stratigraphische und paläofloristische Bedeutung; Habilitationsschrift, Universität Göttingen, 282 p.

1982: Palynological aspects of the Lower/Middle Devonian transition in the Eifel region; Courier Forschungsinstitut Senckenberg, v. 55, p. 279-292.

Riegel, W. and Karathanasopoulos, S.
1982: Palynological criteria for the Siegenian/Emsian transition in the Rhineland; Courier Forschungsinstitut Senckenberg v. 55, p. 199-206.

Rodriguez, R.M.
1978: Mioesporas de la Formación San Pedro/Furada (Silúrico Superior – Devónico Inferior), Cordillera Cantábrica, NO de España; Palinología, Número Extraordinario 1, p. 407-433.

Sandberg, C.A., Streel, M., and Scott, R.A.
1972: Comparison between conodont zonation and spore assemblages at the Devonian-Carboniferous boundary in the western and central United States and in Europe; Compte Rendu, Septième Congrès International de Stratigraphie et de Géologie du Carbonifère, Krefeld, 1971, v. 1, p. 179-203.

Sanford, B.V. and Norris, A.W.
1975: Devonian stratigraphy of the Hudson Platform; Geological Survey of Canada, Memoir 379, 372 p.

Schultz, G.
1968: Eine unterdevonische Mikroflora aus den Klerfer Schichten der Eifel (Rheinisches Schiefergebirge); Palaeontographica, Abteilung B, v. 123, p. 5-42.

Scoffin, T.P.
1971: The conditions of growth of the Wenlock reefs of Shropshire (England); Sedimentology, v. 17, p. 173-219.

Scott, R.A. and Doher, L.I.
1967: Palynological evidence for Devonian age of the Nation River Formation, east-central Alaska; United States Geological Survey Professional Paper 575-B, p. 45-49.

Sennova, V.F.
1972: Spore-pollen assemblages of the Upper Devonian of the Timan-Pechora Province; Geology and Oil and Gas bearing beds of NE European part of USSR, no. 2, Siktivkar, p. 99-107 (in Russian).

Sergeeva, L.A.
1979: Upper Devonian microphytofossils of the Dnepr-Donets Depression; Academy of Sciences of Ukrainian SSR, Institute of Geological Sciences, Kiev, "Naukova Dumka", p. 1-148 (in Russian).

Shepeleva, E.D.
1963: Spores from the Lower Devonian beds of the Podolian Dniester River area; Data on Regional Stratigraphy of USSR, Moscow, p. 98-101 (in Russian).

Shergold, J.H. and Bassett, M.G.
1970: Facies and faunas at the Wenlock/Ludlow boundary of Wenlock Edge, Shropshire; Lethaia, v. 3, no. 2, p. 113-142.

Sleeman, A.G., Higgs, K. and Sevastopulo, G.D.
1983: The stratigraphy of the late Devonian – early Carboniferous rocks of south County Wexford; Geological Survey of Ireland, Bulletin, v. 3, no. 2, p. 141-158.

Sleeman, A.G., Reilly, T.A., and Higgs, K.
1978: Preliminary stratigraphy and palynology of five sections through the Old Head Sandstone and Kinsale formations (Upper Devonian to Lower Carboniferous), on the west side of Cork Harbour; Geological Suvey of Ireland, Bulletin, v. 2, no. 3, p. 167-186.

Smith, D.G.
1975: Wenlock plant spores and tetrads from County Mayo, Ireland; Geological Magazine, v. 112, p. 411-414.

Sorokina, N.L.
1968: Miospore assemblages of Devonian and lower Tournaisian deposits of the southeast of the Dnepr-Donets Depression; Geological Journal, v. 28, no. 4, p. 57-64 (in Russian).

Stanley, E.A.
1969: The occurrence and distribution of pollen and spores in marine sediments; Proceedings of First International Conference on Planktonic Microfossils, Geneva, 1967, v. 2, p. 640-643.

Staplin, F.L.
1961: New plant spores similar to *Torispora* Balme; Journal of Paleontology, v. 35, no. 6, p. 1227-1231.

Steemans, Ph.
1981: Etude stratigraphique des spores dans les couches de transition "Gedinnien-Siegenien" à Nonceveux et à Spa (Belgique); Annales de la Société Géologique de Belgique, v. 104, no. 1, p. 41-59.

1982a: L'âge du poudingue de Fépin (base du Gedinnien) à Lahonry (Belgique); Bulletin de la Société belge de Géologie, v. 90, no. 4, p. 331-340.

Steemans, Ph. (cont.)
1982b: Gedinnian and Siegenian spore stratigraphy in Belgium; Courier Forschungsinstitut Senckenberg, v. 55, p. 165-180.

Streel, M.
1964: Une association de spores du Givétien inférieur de la Vesdre, à Goé (Belgique); Annales de la Société Géologique de Belgique, v. 87, no. 7, p. 1-30.

1965: Etude palynologique du Dévonien du sondage de Booischot (Belgique); Société belge de Géologie, de Paléontologie et d'Hydrologie, Bulletin, v. 73, no. 2, p. 172-185.

1966: Critères palynologiques pour une stratigraphie détaillée du Tnla des bassins Ardenno-Rhénans; Annales de la Société Géologique de Belgique, v. 89, no. 1-4, p. 67-96.

1967: Associations de spores du Dévonien inferieur belge et leur signification stratigraphique; Annales de la Société Géologique de Belgique, v. 90, no. 1-3, p. 11-53.

1972: Dispersed spores associated with *Leclercqia complexa* Banks, Bonamo and Grierson from the late Middle Devonian of Eastern New York State (U.S.A.); Review of Palaeobotany and Palynology, v. 14, p. 205-215.

1977: Corrélations palynologiques dans le Tournaisien du Synclinorium de Namur; Bulletin de la Société Belge de Géologie, v. 82, p. 397-415.

Streel, M., Bless, M., Bouckaert, J., Coen, M., Coen-Aubert, M., Conil, R., Dreesen, R., Dusar, M., Mouravieff, N., and Thorez, J.
1975: Chief micropaleontological limits in the Belgian Upper-Devonian; International Symposium on Belgian Micropaleontological Limits, "Namur 1974", Publication no. 19, p. 1-35.

Streel, M., Demaret-Fairon, M., and Otazo, N.
1975: Siegenian and Emsian spores from the Dinant Basin (Belgium); Commission Internationale de Microflore du Paléozoïque, Newsletter 10, p. 7.

Streel, M., Fairon-Demaret, M., Otazo-Bozo, N., and Steemans, Ph.
1981: Etudes stratigraphiques des spores du Dévonien inférieur au bord sud du synclinorium de Dinant (Belgique) et leurs applications; Annales de la Société Géologique de Belgique, v. 104, no. 1, p. 173-191.

Streel, M. and Traverse, A.
1978: Spores from the Devonian/Mississippian transition near the Horseshoe Curve section, Altoona, Pennsylvania, U.S.A.; Review of Palaeobotany and Palynology, v. 26, no. 1-4, p. 21-39.

Strother, P.K. and Traverse, A.
1979: Plant microfossils from Llandoverian and Wenlockian rocks of Pennsylvania; Palynology, v. 3, p. 1-21.

Sullivan, H.J.
1968: A Tournaisian spore flora from the Cementstone Group of Ayrshire, Scotland; Palaeontology, v. 11, pt. 1, p. 116-131.

Tassonyi, E.J.
1969: Subsurface geology, lower Mackenzie River and Anderson River area, District of Mackenzie; Geological Survey of Canada, Paper 68-25, 207 p.

Taugourdeau-Lantz, J.
1967: Spores nouvelles du Frasnien du Bas Boulonnais (France); Revue de Micropaléontologie v. 10, no. 1, p. 48-60.

1971: Les spores du Frasnien d'une région privilégiée, le Boulonnais; Mémoires de la Société Géologique de France, Mémoire no. 114, p. 1-88.

Tesmer, I.H.,
1963: Geology of Chautauqua County, New York; Pt. 1, Stratigraphy and Paleontology (Upper Devonian); New York State Museum and Science Service, Bulletin 391, p. 1-65.

Teichert, C.
1958: Biostratigraphic concepts; Geological Society of America, Bulletin 69, p. 99-119.

Thorsteinsson, R.
1958: Cornwallis and Little Cornwallis Islands, District of Franklin, Northwest Territories; Geological Survey of Canada, Memoir 294, 134 p.

Thorsteinsson, R. and Uyeno, T.T.
1980: Stratigraphy and conodonts of Upper Silurian and Lower Devonian rocks in the environs of the Boothia Uplift, Canadian Arctic Archipelago; Geological Survey of Canada, Bulletin 292, 75 p.

Timofeev, B.V.
1966: Micropaleophytological study of ancient strata; Academy of Sciences of the USSR, Laboratory of Precambrian Geology, "Nauka", Moscow, 147 p. (in Russian).

Tozer, E.T. and Thorsteinsson, R.
1964: Western Queen Elizabeth Islands, Arctic Archipelago; Geological Survey of Canada, Memoir 332, 242 p.

Tschudy, R.H.
1969: Relationship of palynomorphs to sedimentation; in Aspects of Palynology, R.H. Tschudy and R.A. Scott (eds.); Wiley-Interscience, p. 79-96.

Turnau, E.
1974: Microflora from core samples of some Paleozoic sediments from beneath the flysch Carpathians (Bielsko-Wadowice area, southern Poland); Annales de la Société Géologique de Pologne, v. 44, no. 2-3, p. 143-169.

1975: Microflora of the Famennian and Tournaisian deposits from boreholes of northern Poland; Acta Geologica Polonica, v. 25, no. 4, p. 505-528.

1978: Spore zonation of uppermost Devonian and Lower Carboniferous deposits of western Pomerania; Mededelingen Rijks Geologische Dienst, v. 30-1, p. 1-35.

1979: Korelacje utworów Górnego Dewonu i Karbonu Pomorza Zachodniego w oparciu o badania miosporowe; Rocznik Polskiego Towarzystwa Geologicznego, Kraków, v. 49, no. 3-4, p. 231-269.

Van der Zwan, C.J.
1979: Aspects of Late Devonian and Early Carboniferous palynology of southern Ireland. I. The *Cyrtospora cristifer* morphon; Review of Palaeobotany and Palynology, v. 28, no. 1, p. 1-20.

1980: Aspects of Late Devonian and Early Carboniferous palynology of southern Ireland. III. Palynology of Devonian-Carboniferous transition sequences with special reference to the Bantry Bay area, Co. Cork; Review of Palaeobotany and Palynology, v. 30, no. 3-4, p. 165-286.

Van der Zwan, C.J. and Walton, H.S.
1981: The *Cyrtospora cristifer* morphon: inclusion of *Cornispora varicornata* and *C. monocornata*; Review of Palaeobotany and Palynology, v. 33, p. 139-152.

Van Veen, P.M.
1981: Aspects of Late Devonian and Early Carboniferous palynology of southern Ireland. IV. Morphological variation within *Diducites*, a new form genus to accommodate camerate spores with two-layered outer wall; Review of Palaeobotany and Palynology, v. 31, p. 261-287.

Vavrdova, M.
1982: Recycled acritarchs in the uppermost Ordovician of Bohemia; Casopis pro Mineralogii a Geologii, v. 27, no. 4, p. 337-345.

Vigran, J.O.
1964: Spores from Devonian deposits, Mimerdalen, Spitsbergen; Norsk Polarinstitutt, Skr. 132, p. 1-32.

Visscher, H.
1971: The Permian and Triassic of the Kingscourt Outlier, Ireland; Geological Survey of Ireland, Special Paper no. 1, 114 p.

Volochaev, F.Ya., Kud'yarov, I.S. and Petrenko, V.I.
1979: Age of the volcanogenic-sedimentary formation overlying alluvial bauxite in Middle Timan; Proceedings of the USSR Academy of Sciences, Geological Series, v. 1979, no. 5, p. 151-154 (in Russian).

Wang Yu, Yu Chang-ming, Xu Han-kui, Liao Wei-hua, and Tsai Chung-yang
1978: Devonian biostratigraphy of South China; Nanking Institute of Geology and Palaeontology, Papers for the International Symposium on the Devonian System, no. 1, p. 1-9.

Weddige, K.
1977: Die Conodonten der Eifel-Stufe im Typusgebiet und in benachbarten Fazies gebieten; Senckenbergiana Lethaea, v. 58, p. 271-419.

Weddige, K., Werner, R. and Ziegler, W.
1979: The Emsian-Eifelian boundary, an attempt at correlation between the Eifel and Ardennes regions; Newsletters on Stratigraphy, v. 8, no. 2, p. 159-169.

White, E.I.
1950: The vertebrate faunas of the Lower Old Red Sandstone of the Welsh borders; Bulletin of the British Museum (Natural History), Geology Series, v. 1, no. 3, p. 51-67.

Whiteley, M.J.
1980: Givetian and Frasnian spores from the Key Point well, Parry Islands, Arctic Canada; Review of Palaeobotany and Palynology, v. 29, p. 301-311.

Williams, B.P.J. (ed.)
1978: Excursion B2. The Old Red Sandstone of the Welsh Borderland and South Wales, field guide; in A field guide to selected outcrop areas of the Devonian of Scotland, the Welsh Borderland and South Wales; Palaeontological Association, International Symposium on the Devonian System, p. 55-106.

Winslow, M.R.
1962: Plant spores and other microfossils from Upper Devonian and Lower Mississippian rocks of Ohio; United States Geological Survey, Professional Paper 364, 93 p.

Wood, G.D.
1978: Palynology and paleobotany of the Java and lowermost Canadaway formations, Upper Devonian (Senecan/Chautauquan) New York State; Michigan State University, unpublished doctoral thesis, 241 p.

Ziegler, W.
1979: Historical subdivisions of the Devonian; in The Devonian System, M.R. House, C.T. Scrutton, and M.G. Bassett (eds.); Palaeontological Association, Special Paper 23, p. 23-47.

Ziegler, W. and Klapper, G.
1982: The *disparilis* conodont zone, the proposed level for the Middle-Upper Devonian boundary; Courier Forschungsinstitut Senckenberg, v. 55, p. 463-492.

Ziegler, W. and Sandberg, C.A.
1984: *Palmatolepis*-based revision of upper part of standard Late Devonian conodont zonation; in Conodont biofacies and provincialism, D.L. Clark (ed.); Geological Society of America, Special Paper 196, p. 179-194.

## APPENDIX

### Locality register for specimens figured herein, for which descriptions of localities have not been published

Plate 1, figure 4. Pennsylvania State University "Mill Hall locality", 1.1 m southeast of Mill Hall, Clinton County, Pennsylvania; outcrop at a level 250 m above US Route 220.

Plate 11, figures 5 and 6. GSC locality C-84874. Southwestern Ellesmere Island, District of Franklin, NTS 49C, UTM zone 16, 519550E, 8569100N, east of Bird Fiord; outcrop 8 m above base of Strathcona Fiord Formation.

Plate 14, figure 13. GSC plant locality 7013. Northwestern Cornwallis Island, District of Franklin, NTS 58G/5; outcrop on Rookery Creek, Locality 32 ("Pennsylvanian") of Thorsteinsson (1958, p. 111).

Plate 15, figure 6. GSC locality C-84874 (same as Plate 11, figures 5 and 6).

Plate 15, figure 7. GSC locality C-28109. Western District of Mackenzie, NTS 96E/11, 65°30'20"N, 127°13'26"W, Banff Aquit GPD Oscar Creek H-71 well; cuttings from 640-730 ft (195-222.5 m), spore-bearing member of Hare Indian Formation (see Tassonyi, 1969).

Plate 16, figure 4. GSC locality C-33961. Banks Island, District of Franklin, NTS 98A/16, 72°47'51"N, 120°44'48"W, Elf Texaco Tiritchik M-48 well; cuttings from 3100-3140 ft (945-957 m), 884-924 ft (269.5-282 m) above base of Weatherall Formation.

Plate 16, figure 5. GSC locality C-91916. Southwestern Ellesmere Island, District of Franklin, NTS 49C/6, UTM zone 16, 505250E, 8562675N; outcrop on east side of inner part of southeast arm of Bird Fiord; Fram Formation.

Plate 16, figures 6 and 7. GSC locality C-91913 (same as Plate 16, figure 5, but 50.5 m lower stratigraphically).

Plate 17, figure 6. GSC locality C-91918 (same as Plate 16, figure 5, but 17.9 m higher stratigraphically).

Plate 17, figure 7. GSC locality 97081. East-central British Columbia, NTS 93H/16, Monkman Pass area, 53°52'34"N, 120°01'08"W; outcrop 19 m below top of Perdrix Formation.

Plate 18, figure 6. GSC locality D1726. Antigonish County, Nova Scotia, NTS 11F/12, 45°44'24.4"N, 61°53'28.46"W, west side of St. George's Bay, north of McIsaac Point.

Plate 18, figure 7. GSC plant locality 5436. Northwestern District of Mackenzie, NTS 106M, about 48 km southwest of Fort McPherson; outcrop along Vittrekwa River, Imperial Formation.

Plate 18, figure 9. Richardson locality NY96/1965. McKean County, Pennsylvania; road cut on U.S. Route 6 at Bush Hill, 5 km east of Smethport.

Plate 19, figure 1. GSC plant locality 5436 (same as Plate 18, figure 7).

Plate 19, figure 5. Richardson locality US6/1968. Pope Hollow section, along boundary between Caroll and South Valley townships, Jamestown Quadrangle near Chautauqua, New York.

Plate 19, figure 9. Richardson locality PA19/1966. Warren County, Pennsylvania, road cut on State Highway 59, 19.3 km east of Warren and 2.25 km northeast of Kinzua Dam.

Plate 20, figure 1. Richardson locality NY95/1965. McKean County, Pennsylvania, Hanley Quarries, east of Lewis Run.

Plate 20, figure 3 (same as Plate 19, figure 5).

Plate 20, figure 5. Richardson locality NY25/1965. Cattaraugus County, New York State, road cut on State Highway 16A, 4.8 km south of Olean.

Plate 20, figure 7. GSC locality D1726 (same as Plate 18, figure 6).

Plate 21, figure 4. GSC plant locality 6496. Yukon Territory, NTS 106L, Caribou River; Imperial Formation.

Plate 21, figure 6. GSC locality D1726 (same as Plate 18, figure 6).

Plate 21, figure 7. GSC locality 97057. Northernmost tip of Western Australia, Amax Ellendale well, 1750.5 m; Fairfield Formation(?).

## PLATES 1-21

Illustrations of spores

Letter-number designations following descriptions of figures are type specimen numbers. For further explanation, see Table 2 (in pocket)

## PLATE 1

(All figures are X1000, unless stated otherwise)

Figure 1. *Ambitisporites avitus* Hoffmeister. FM 1.

Figure 2. *Ambitisporites dilutus* (Hoffmeister) Richardson and Lister. FM 2.

Figure 3. *Retusotriletes warringtonii* Richardson and Lister. FM 3.

Figures 4, 5. *Tetrahedraletes medinensis* Strother and Traverse; 4, FM4; 5, Scanning Electron Micrograph, X2000.

Figures 6-8. *Archaeozonotriletes chulus* (Cramer) Richardson and Lister var. *chulus*; 6, 7, FM 5; 8, two specimens, one laterally compressed, FM 6.

Figure 9. *Archaeozonotriletes chulus* (Cramer) Richardson and Lister var. *nanus* Richardson and Lister. GSC 41705.

Figure 10. ?*Archaeozonotriletes* cf. *divellomedium* Chibrikova. FM 8.

Figures 11, 13, 14. cf. *Synorisporites verrucatus* Richardson and Ioannides; 11, X1500, FM 12; 13, 14, FM 13.

Figure 12. *Emphanisporites* cf. *protophanus* Richardson and Ioannides. FM 11.

Figures 15, 16. *Retusotriletes abundo* Rodriguez; 15, FM 9; 16, FM 10.



1

2





3

4



8

6

7



15

9

10



11



12

13



14



16

## PLATE 2

(All figures are X1000, unless stated otherwise)

Figures 1-3. *Synorisporites libycus* Richardson and Ioannides; 1, X2000, FM 14; 2, 3, FM 15.

Figure 4. *Emphanisporites neglectus* Vigran. FM 19.

Figures 5, 10. *Synorisporites* cf. *verrucatus* Richardson and Ioannides; 5, FM 17; 10, X1500, FM 16.

Figures 6, 7. ?*Lophozonotriletes poecilomorphus* Richardson and Ioannides. FM 21.

Figures 8, 9. cf. *Brochotriletes* sp. A of Richardson and Ioannides, 1973, X1500. FM 22.

Figure 11. cf. *Synorisporites downtonensis* Richardson and Lister. IGS MPK 10.

Figure 12. *Emphanisporites protophanus* Richardson and Ioannides. FM 20.

Figure 13. *Amicosporites splendidus* Cramer. FM 26.

Figure 14. *Apiculiretusispora spicula* Richardson and Lister, holotype. FM 24.

Figure 15. *Apiculiretusispora synorea* Richardson and Lister. FM 25.

Figure 16. *Tetraletes variabilis* Cramer. FM 23.

Figures 17, 18. *Synorisporites tripapillatus* Richardson and Lister, holotype, X2000. FM 18.


1


2


3


4


5


6


7


8

9


10

11

12


13

14

15


16


17


18

## PLATE 3

(All figures are X1000, unless stated otherwise)

Figure 1. *Ambitisporites* sp. B of Richardson and Ioannides, 1973. FM 27.

Figure 2. *Cymbosporites echinatus* Richardson and Lister. FM 28.

Figure 3. *Emphanisporites splendens* Richardson and Ioannides, holotype. FM 30.

Figures 4, 5. *Cymbosporites verrucosus* Richardson and Lister, holotype. FM 29.

Figures 6, 7. *Synorisporites verrucatus* Richardson and Lister, holotype. FM 32.

Figures 8, 9. *Synorisporites downtonensis* Richardson and Lister. FM 31.

Figures 10, 11. *Streelispora newportensis* (Chaloner and Streel) Richardson and Lister, X2000. FM 33.

Figures 12, 13. *Acinosporites salopiensis* Richardson and Lister, holotype, X2000. FM 34.



1



2

3

4



5



6

7

8



9



12



10



11

13

## PLATE 4

(All figures are X1000, unless stated otherwise)

Figure 1. *Emphanisporites micromatus* Richardson and Lister, X2000. FM 35.

Figure 2. *Apiculiretusispora plicata* (Allen) Streel. GSC 15153.

Figures 3, 4. *Cymbosporites proteus* McGregor and Camfield, holotype, X500. GSC 41710.

Figure 5. *Emphanisporites epicautus* Richardson and Lister. FM 37.

Figures 6, 7. *Chelinospora cassicula* Richardson and Lister, holotype. FM 36.

Figures 8, 9. *Perotrilites microbaculatus* Richardson and Lister. FM 38.

Figure 10. *Retusotriletes maculatus* McGregor and Camfield, holotype. GSC 41706.

Figure 11. *Breconisporites breconensis* Richardson, Streel, Hassan and Steemans, holotype, X1100. V61174/FM 39.

Figure 12. *Camptozonotriletes* sp. cf. *C. caperatus* McGregor. FM 40.

Figure 13. *Emphanisporites zavallatus* Richardson, Streel, Hassan and Steemans, holotype. V61170/FM 41.

Figure 14. *Cirratriradites* sp. A of Streel et al., 1981. FM 42.

Figures 15, 16. *Clivosispora verrucata* McGregor var. *verrucata*, holotype, X500. GSC 15179.

1



2



6



3

4



5



7

8

10



11



9



12



13

14



15



16

## PLATE 5

(All figures are X1000, unless stated otherwise)

Figures 1, 2. *Verrucosisporites polygonalis* Lanninger. GSC 31989.

Figure 3. *Brochotriletes foveolatus*? Naumova. GSC 31990.

Figures 4, 5. *Dictyotriletes emsiensis* (Allen) McGregor. GSC 41734.

Figure 6. *Dibolisporites eifeliensis* (Lanninger) McGregor, X500. GSC 49817.

Figures 7, 8. *Dibolisporites wetteldorfensis* Lanninger. GSC 15161.

Figure 9. *Apiculiretusispora* cf. *brandtii* Streel. IGS MPK 1895.

Figures 10, 11. *Clivosispora verrucata* McGregor var. *convoluta* McGregor and Camfield, X500. GSC 27081.

Figures 12, 13. *Emphanisporites decoratus* Allen. GSC 27076.

1



2



3



4



5



6



7



8

10



12

9



11

13

## PLATE 6

(All figures are X1000, unless stated otherwise)

| | | |
|---|---|---|
| Figures | 1, 2. | *Dictyotriletes favosus* McGregor and Camfield, X500. GSC 15181. |
| Figures | 3, 4. | *Brochotriletes hudsonii* McGregor and Camfield, holotype, X500. GSC 41729. |
| Figure | 5. | *Dibolisporites* n. sp. K of Hassan, 1982. FM 43. |
| Figure | 6. | *Apiculiretusispora arenorugosa* McGregor, holotype, X500. GSC 31961. |
| Figure | 7. | *Camptozonotriletes caperatus* McGregor, holotype, X500. GSC 15197. |
| Figure | 8. | *Acinosporites lindlarensis* Riegel var. *lindlarensis*. GSC 73289. |
| Figures | 9, 10. | *Emphanisporites annulatus* McGregor. GSC 32002. |
| Figures | 11, 12. | *Camarozonotriletes sextantii* McGregor and Camfield. GSC 27083. |



1



2



7

8



3

5

9

10



4



6



11



12

## PLATE 7

(All figures are X500, unless stated otherwise)

Figure 1. *Apiculatasporites microconus* (Richardson) McGregor and Camfield. FM 44.

Figures 2, 3. *Dibolisporites echinaceus* (Eisenack) Richardson; 3, X1200. GSC 31972.

Figure 4. *Calyptosporites biornatus* (Lanninger) Richardson.

Figure 5. *Apiculiretusispora brandtii* Streel.

Figure 6. *Dictyotriletes gorgoneus* Cramer, X1000. GSC 31997.

Figure 7. *Camptozonotriletes* cf. *aliquantus* Allen, X1000.

Figure 8. *Dictyotriletes canadensis* McGregor. GSC 32001.

Figure 9. *Emphanisporites schultzii* McGregor, X1000. GSC 32008.

Figure 10. *Verruciretusispora dubia* (Eisenack) Richardson and Rasul. IGS MPK 1897.



1

2

3



4

5

6

7

8

9

10

## PLATE 8

(All figures are X500, unless stated otherwise)

Figures 1, 2. *Ancyrospora eurypterota* Riegel; 2, X1000.

Figure 3. *Grandispora douglastownense* McGregor, holotype. GSC 32034.

Figures 4, 5. *Acinosporites apiculatus* (Streel) Streel; 4, X2000.

Figure 6. *Ancyrospora loganii* McGregor. GSC 15282.

Figures 7, 8. *Ancyrospora nettersheimensis* Riegel; 8, X1000.



1



2

3

4

5

6



7



8

## PLATE 9

(All figures are X500, unless stated otherwise)

Figure 1. *Ancyrospora kedoae* (Riegel) Turnau.

Figure 2. *Apiculiretusispora gaspiensis* McGregor, X1000. GSC 31968.

Figure 3. *Calyptosporites* sp. nov. of Riegel, 1975.

Figure 4. *Calyptosporites proteus* (Naumova) Allen. GSC 49834.

Figure 5. *Grandispora eximia* (Allen) McGregor and Camfield. GSC 66310.

Figure 6. *Spinozonotriletes arduinnae* Riegel, holotype.

Figure 7. *Hystricosporites* cf. *H. corystus* Richardson.

1

2



3

4



5

6

7

## PLATE 10

(All figures are X500, unless stated otherwise)

Figures 1, 2. *Hystricosporites microancyreus* Riegel.

Figures 3, 4. *Acinosporites acanthomammillatus* Richardson, holotype. FM 45.

Figure 5. *Rhabdosporites langii* (Eisenack) Richardson. FM 46.

Figure 6. *Calyptosporites velatus* (Eisenack) Richardson. FM 47.

Figure 7. *Anapiculatisporites petilus* Richardson, holotype, X1000. FM 48.

Figure 8. *Kraeuselisporites olli*? (Chibrikova) McGregor and Camfield. GSC 29391.

Figure 9. *Acinosporites macrospinosus* Richardson, holotype. FM 49.



1



2

3



4



5



6

7

8

9

## PLATE 11

(All figures are X500)

Figures 1, 2. *Ancyrospora ancyrea* (Eisenack) Richardson var. *ancyrea*. FM 50.

Figures 3, 4. *Kraeuselisporites acerosus* (Arkhangelskaya) McGregor and Camfield. GSC 66349.

Figures 5, 6. *Periplecotriletes tortus* Egorova. GSC 73290.

Figures 7, 8. *Grandispora longa* Chi and Hills. GSC 15305.



1

3

5

4

6

2



8

7

## PLATE 12

(All figures are X500, unless stated otherwise)

Figure 1. *Corystisporites multispinosus* Richardson, holotype. FM 51.

Figure 2. *Densosporites concinnus* (Owens) McGregor and Camfield, X1000. GSC 73291.

Figure 3. *Perotrilites bifurcatus* Richardson, holotype. FM 52.

Figure 4. *Retusotriletes rugulatus* Riegel, X1000. GSC 66379.

Figure 5. *Retusotriletes distinctus* Richardson. FM 53.

Figure 6. *Densosporites devonicus* Richardson. FM 54.

Figure 7. *Densosporites inaequus* (McGregor) McGregor and Camfield. GSC 66274.

Figure 8. *Grandispora mammillata* Owens. GSC 66320.



1 2

3 4 5



6 7 8

## PLATE 13

(All figures are X500, unless stated otherwise)

Figure 1. *Grandispora naumovii* (Kedo) McGregor. FM 55.

Figure 2. *Cristatisporites orcadensis* Richardson, holotype. FM 56.

Figure 3. *Lophotriletes devonicus* (Naumova ex Chibrikova) McGregor and Camfield, X1000. GSC 66356.

Figure 4. *Perotrilites conatus* Richardson. FM 57.

Figure 5. *Grandispora inculta* Allen, X1000. GSC 66317.

Figure 6. *Kraeuselisporites rugosus* (Owens) McGregor and Camfield. GSC 29387.

Figure 7. *Ancyrospora grandispinosa* Richardson, X300. FM 58.



1



2



3

4

5



6



7

## PLATE 14

(All figures are X500, unless stated otherwise)

Figure 1. *Hystricosporites gravis* Owens. GSC 15550.

Figure 2. *Verrucosisporites scurrus* (Naumova) McGregor and Camfield, X1000. GSC 66396.

Figures 3, 4. *Retispora archaelepidophyta* (Kedo) McGregor and Camfield; 3, GSC 49843; 4, X1000, GSC 66372.

Fiugres 5, 6. *Verrucosisporites premnus* Richardson; 6, X1000. GSC 66394.

Figure 7. *Hystricosporites reflexus* Owens, holotype. GSC 15546.

Figures 8, 9. *Geminospora lemurata* Balme, X1000. GSC 66296.

Figure 10. *Archaeozonotriletes variabilis* Naumova. GSC 13624.

Figure 11. *Cymbosporites magnificus* (McGregor) McGregor and Camfield, holotype. GSC 13036.

1

2



3



4



5



6

7

8

9

10



11

## PLATE 15

(All figures are X500, unless stated otherwise)

Figure 1. *Biharisporites parviornatus* Richardson, holotype. FM 59.

Figure 2. *Geminospora? bislimbata* (Chibrikova) McGregor and Camfield. GSC 66295.

Figure 3. *Perotrilites heclaensis* McGregor and Camfield, X1000. GSC 66362.

Figures 4, 5. *Archaeozonotriletes timanicus* Naumova. GSC 66216.

Figure 6. *Corystisporites serratus* (Naumova) McGregor and Camfield, X1000. GSC 73292.

Figure 7. *Hymenozonotriletes celeber* Chibrikova. GSC 49845.

Figure 8. *Ancyrospora involucra* Owens. GSC 15684.

Figure 9. *Cristatisporites triangulatus* (Allen) McGregor and Camfield. GSC 49842.

Figure 10. *Aneurospora greggsii* (McGregor) Streel, holotype. GSC 13184.



1

2

3



4

5



6

7



8

9

10

## PLATE 16

(All figures are X500, unless stated otherwise)

Figure 1. *Contagisporites optivus* (Chibrikova) Owens var. *optivus*. GSC 49848.

Figures 2, 3. *Ancyrospora langii* (Taugourdeau-Lantz) Allen. GSC 73293.

Figure 4. *Aneurospora goensis* Streel. GSC 49838.

Figure 5. *Chelinospora ligurata* Allen. GSC 73294.

Figures 6, 7. *Ancyrospora melvillensis* Owens. GSC 73295.

Figure 8. *Grandispora tomentosa* Taugourdeau-Lantz. GSC 41770.

Figure 9. *Archaeoperisaccus timanicus* Pashkevich. GSC 29412.

Figure 10. *Chelinospora concinna* Allen, X1000. GSC 49839.

1



2

3

4

5



6



7



8



9



10

## PLATE 17

(All figures are X500, unless stated otherwise)

Figure 1. *Archaeoperisaccus ovalis* Naumova, X1000. GSC 73296.

Figure 2. *Verrucosisporites bulliferus* nom. nov., X1000. GSC 73297.

Figure 3. *Archaeoperisaccus opiparus* Owens. GSC 73298.

Figures 4, 5. *Ancyrospora furcula* Owens; 5, X1000. GSC 49847.

Figure 6. *Hymenozonotriletes deliquescens* Naumova. GSC 73299.

Figure 7. cf. *Archaeozonotriletes laetus* Sergeeva, X1000. GSC 73300.

Figures 8, 9. *Nikitinsporites canadensis* Chaloner; 8, GSC 73301; 9, X150, GSC 43732.

Figure 10. *Lophozonotriletes media* Taugourdeau-Lantz. GSC 73302.

Figure 11. *Diaphanospora reticulata* Guennel, X1000. GSC 73303.

1



2



3

4



6



7



5

8



10

9



11

## PLATE 18

(All figures are X1000, unless stated otherwise)

Figure 1. *Auroraspora torquata* Higgs, holotype, X500.

Figures 2, 3. *Grandispora gracilis* (Kedo) Streel; 3, X5000.

Figure 4. *Archaeozonotriletes famenensis* Naumova. GSC 49851.

Figure 5. *Cornispora varicornata* Staplin and Jansonius. GSC 49852.

Figure 6. *Auroraspora hyalina* (Naumova) Streel. GSC 73304.

Figure 7. *Cyrtospora cristifera* (Luber) Van der Zwan. GSC 49853.

Figure 8. *Auroraspora pseudocrista* Ahmed. FM 60.

Figure 9. *Diducites versabilis* (Kedo) Van Veen. FM 67.

Figure 10. *Hystricosporites multifurcatus* (Winslow) Mortimer and Chaloner, X500. GSC 73305.

1



2

3

4

5

6



7



8

9

10

## PLATE 19

(All figures are X1000, unless stated otherwise)

Figure 1. *Knoxisporites dedaleus* (Naumova) Streel. GSC 73306.

Figures 2, 3. "(?)*Hymenozonotriletes denticulatus* Naumova" of McGregor, 1967. GSC 49850.

Figure 4. *Diducites poljessicus* (Kedo) Van Veen.

Figure 5. *Crassispora catenata* Higgs. FM 62.

Figure 6. *Lophozonotriletes lebedianensis* Naumova. GSC 15422.

Figure 7. *Rugospora flexuosa* (Jushko) Streel. GSC 73308.

Figure 8. *Auroraspora macra* Sullivan. GSC 73309.

Figure 9. *Lagenicula* sp. A of Streel in Becker et al., 1974, X500. FM 61.

1

2

3

4



5



6



7



8

9

## PLATE 20

(All figures are X1000, unless stated otherwise)

Figure 1. *Grandispora cornuta* Higgs. FM 63.

Figure 2. *Retispora cassicula* (Higgs) Higgs and Russell, X500.

Figure 3. *Retusotriletes phillipsii* Clendening, Eames and Wood. FM 64.

Figure 4. *Retispora lepidophyta* (Kedo) Playford. GSC 73310.

Figure 5. *Vallatisporites pusillites* (Kedo) Dolby and Neves. FM 65.

Figure 6. *Grandispora echinata* Hacquebard. GSC 25496.

Figure 7. *Discernisporites micromanifestus* (Hacquebard) Sabry and Neves. GSC 73311.



1



2

3

4

5



6



7

## PLATE 21

(All figures are X1000, unless stated otherwise)

Figure 1. *Raistrickia variabilis* Dolby and Neves. GSC 73312.

Figures 2, 3. *Tumulispora rarituberculata* (Luber) Turnau. FM 66.

Figure 4. *Tumulispora malevkensis* (Kedo) Turnau. GSC 49854.

Figure 5. *Knoxisporites literatus* (Waltz) Playford. GSC 25450.

Figure 6. *Hymenozonotriletes explanatus* (Luber) Kedo, X500. GSC 73313.

Figure 7. *Verrucosisporites nitidus* (Naumova) Playford. GSC 73314.

Figure 8. *Vallatisporites verrucosus* Hacquebard, X500.

Figure 9. *Raistrickia corynoges* Sullivan. GSC 73315.

PLATE 21



1

2

1

3

6

4

5



7

8

9

MAPS NOT

FILMED

CARTES

OMISES